

S-100A3 (T-12): sc-132084

BACKGROUND

The S-100 protein family consists of a group of calcium-binding proteins that are exclusively expressed in vertebrates and exhibit cell and tissue-specific expression. The expression levels of its members differ in various pathological conditions. The extracellular functions of the S-100 family may include the ability to enhance neurite outgrowth, involvement in inflammation and motility of tumor cells. S-100A3 (S100 calcium binding protein A3), also known as S100E, is a 101 amino acid cysteine-rich protein that belongs to the S-100 family of proteins. Expressed specifically at the inner endocuticle of hair fibers, S-100A3 contains two EF-hand domains and exists as a multimer. S-100A3 binds calcium with low affinity and zinc with high affinity and is believed to play a role in the calcium-dependent differentiation of cuticle cells and in the formation of hair shafts.

REFERENCES

- Engelkamp, D., et al. 1993. Six S100 genes are clustered on human chromosome 1q21: identification of two genes coding for the two previously unreported calcium-binding proteins S-100D and S-100E. *Proc. Natl. Acad. Sci. USA* 90: 6547-6551.
- Schäfer, B.W., et al. 1995. Isolation of a YAC clone covering a cluster of nine S-100 genes on human chromosome 1q21: rationale for a new nomenclature of the S-100 calcium-binding protein family. *Genomics* 25: 638-643.
- Mittl, P.R., et al. 2002. Metal-free MIRAS phasing: structure of apo-S-100A3. *Acta Crystallogr. D Biol. Crystallogr.* 58: 1255-1261.
- Kizawa, K., et al. 2002. Characterization of the cysteine-rich calcium-binding S-100A3 protein from human hair cuticles. *Biochem. Biophys. Res. Commun.* 299: 857-862.
- Fritz, G., et al. 2002. The crystal structure of metal-free human EF-hand protein S-100A3 at 1.7-Å resolution. *J. Biol. Chem.* 277: 33092-33098.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 176992. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Kizawa, K., et al. 2005. Dissimilar effect of perming and bleaching treatments on cuticles: advanced hair damage model based on elution and oxidation of S-100A3 protein. *J. Cosmet. Sci.* 56: 219-226.
- Heizmann, C.W., et al. 2007. Pathologies involving the S-100 proteins and RAGE. *Subcell. Biochem.* 45: 93-138.
- Kizawa, K., et al. 2008. Specific citrullination causes assembly of a globular S-100A3 homotetramer: a putative Ca²⁺ modulator matures human hair cuticle. *J. Biol. Chem.* 283: 5004-5013.

CHROMOSOMAL LOCATION

Genetic locus: S100A3 (human) mapping to 1q21.3; S100a3 (mouse) mapping to 3 F1.

SOURCE

S-100A3 (T-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of S-100A3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-132084 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

S-100A3 (T-12) is recommended for detection of S-100A3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other S-100 family members.

S-100A3 (T-12) is also recommended for detection of S-100A3 in additional species, including canine and porcine.

Suitable for use as control antibody for S-100A3 siRNA (h): sc-78754, S-100A3 siRNA (m): sc-153188, S-100A3 shRNA Plasmid (h): sc-78754-SH, S-100A3 shRNA Plasmid (m): sc-153188-SH, S-100A3 shRNA (h) Lentiviral Particles: sc-78754-V and S-100A3 shRNA (m) Lentiviral Particles: sc-153188-V.

Molecular Weight of S-100A3 monomer: 12 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfation
Guaranteed

Try **S-100A3 (A-11): sc-514339**, our highly recommended monoclonal alternative to S-100A3 (T-12).