

TRIM36 (C-13): sc-132127

BACKGROUND

TRIM36 (tripartite motif-containing 36), also known as RNF98 (RING finger protein 98), HAPRIN (haploid germ cell-specific RBCC protein) or RBCC728, is a 728 amino acid protein that belongs to the TRIM/RBCC (RING finger, B box, coiled-coil) family. Predominantly expressed in prostate, testis and brain with weak expression in heart, kidney and lung, TRIM36 contains two B box-type zinc fingers, a SPRY domain, a coiled-coil domain, a fibronectin type-III domain and a RING-type zinc finger; a motif that has zinc-chelating activity and is involved in mediating protein-protein and protein-DNA interactions. Localizing to the cytoplasm and the acrosomal region of germ cells and mature sperm, TRIM36 is believed to play a role in the acrosome reaction and fertilization. In addition, TRIM36 is overexpressed in prostate cancer, suggesting a possible role for TRIM36 in prostate tumorigenesis.

REFERENCES

1. Reymond, A., Meroni, G., Fantozzi, A., Merla, G., Cairo, S., Luzi, L., Riganelli, D., Zanaria, E., Messali, S., Cainarca, S., Guffanti, A., Minucci, S., Pelicci, P.G. and Ballabio, A. 2001. The tripartite motif family identifies cell compartments. *EMBO J.* 20: 2140-2151.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 609317. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Kitamura, K., Tanaka, H. and Nishimune, Y. 2003. Haprin, a novel haploid germ cell-specific RING finger protein involved in the acrosome reaction. *J. Biol. Chem.* 278: 44417-44423.
4. Balint, I., Müller, A., Nagy, A. and Kovacs, G. 2004. Cloning and characterization of the RBCC728/TRIM36 zinc-binding protein from the tumor suppressor gene region at chromosome 5q22.3. *Gene* 332: 45-50.
5. Kitamura, K., Tanaka, H. and Nishimune, Y. 2005. The RING-finger protein haprin: domains and function in the acrosome reaction. *Curr. Protein Pept. Sci.* 6: 567-574.
6. Kitamura, K., Nishimura, H., Nishimune, Y. and Tanaka, H. 2005. Identification of human HAPRIN potentially involved in the acrosome reaction. *J. Androl.* 26: 511-518.
7. Short, K.M. and Cox, T.C. 2006. Subclassification of the RBCC/TRIM superfamily reveals a novel motif necessary for microtubule binding. *J. Biol. Chem.* 281: 8970-8980.

CHROMOSOMAL LOCATION

Genetic locus: TRIM36 (human) mapping to 5q22.3; Trim36 (mouse) mapping to 18 C.

SOURCE

TRIM36 (C-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of TRIM36 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-132127 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

TRIM36 (C-13) is recommended for detection of TRIM36 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other TRIM36 family members.

TRIM36 (C-13) is also recommended for detection of TRIM36 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for TRIM36 siRNA (h): sc-91989, TRIM36 siRNA (m): sc-154647, TRIM36 shRNA Plasmid (h): sc-91989-SH, TRIM36 shRNA Plasmid (m): sc-154647-SH, TRIM36 shRNA (h) Lentiviral Particles: sc-91989-V and TRIM36 shRNA (m) Lentiviral Particles: sc-154647-V.

Molecular Weight of TRIM36: 83 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfaction
 Guaranteed

Try **TRIM36 (SS-03): sc-100881**, our highly recommended monoclonal alternative to TRIM36 (C-13).