

CELF3 (P-12): sc-132225

BACKGROUND

CELF3 (CUG-BP- and ETR-3-like factor 3), also known as TNRC4 (trinucleotide repeat containing 4), CAGH4 (CAG repeat protein 4), ETR-1 (ELAV-type RNA-binding protein 1), ERDA4 (expanded repeat domain protein CAG/CTG 4) or BRUNOL1 (Bruno-like protein 1), is an RNA-binding protein that belongs to the CELF/BRUNOL family of proteins. Expressed specifically in brain, CELF3 contains three RRM (RNA recognition motif) domains (two at the N-terminus and one at the C-terminus) and localizes to the nucleus and the cytoplasm. As is characteristic of the CELF/BRUNOL family, CELF3 binds specifically to muscle-specific splicing enhancer (MSE)-containing RNAs and is believed to play a role in the regulation of pre-mRNA alternative splicing. In addition, CELF3 may participate in translation and mRNA editing. Due to alternative splicing events, CELF3 exists as three isoforms.

REFERENCES

- Margolis, R.L., Abraham, M.R., Gatchell, S.B., Li, S.H., Kidwai, A.S., Breschel, T.S., Stine, O.C., Callahan, C., McInnis, M.G. and Ross, C.A. 1997. cDNAs with long CAG trinucleotide repeats from human brain. *Hum. Genet.* 100: 114-122.
- Good, P.J., Chen, Q., Warner, S.J. and Herring, D.C. 2000. A family of human RNA-binding proteins related to the *Drosophila* Bruno translational regulator. *J. Biol. Chem.* 275: 28583-28592.
- Ladd, A.N., Charlet, N. and Cooper, T.A. 2001. The CELF family of RNA binding proteins is implicated in cell-specific and developmentally regulated alternative splicing. *Mol. Cell. Biol.* 21: 1285-1296.
- Aldenhoven, J., Chen, Y., Backofen, B. and Moran, C. 2003. Improving the comparative map of porcine chromosome 10 with respect to human chromosomes 1, 9 and 10. *Cytogenet. Genome Res.* 102: 121-127.
- Ladd, A.N., Nguyen, N.H., Malhotra, K. and Cooper, T.A. 2004. CELF6, a member of the CELF family of RNA-binding proteins, regulates muscle-specific splicing enhancer-dependent alternative splicing. *J. Biol. Chem.* 279: 17756-17764.
- Singh, G., Charlet-B, N., Han, J. and Cooper, T.A. 2004. ETR-3 and CELF4 protein domains required for RNA binding and splicing activity *in vivo*. *Nucleic Acids Res.* 32: 1232-1241.
- Chapple, J.P., Anthony, K., Martin, T.R., Dev, A., Cooper, T.A. and Gallo, J.M. 2007. Expression, localization and tau exon 10 splicing activity of the brain RNA-binding protein TNRC4. *Hum. Mol. Genet.* 16: 2760-2769.
- Dev, A., Nayernia, K., Meins, M., Adham, I., Lacone, F. and Engel, W. 2007. Mice deficient for RNA-binding protein brunol1 show reduction of spermatogenesis but are fertile. *Mol. Reprod. Dev.* 74: 1456-1464.

CHROMOSOMAL LOCATION

Genetic locus: TNRC4 (human) mapping to 1q21.3; Tnrc4 (mouse) mapping to 3 F2.1.

SOURCE

CELF3 (P-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CELF3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-132225 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CELF3 (P-12) is recommended for detection of CELF3 isoforms 1-3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with CELF5 or CELF6.

CELF3 (P-12) is also recommended for detection of CELF3 isoforms 1-3 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for CELF3 siRNA (h): sc-88405, CELF3 siRNA (m): sc-142261, CELF3 shRNA Plasmid (h): sc-88405-SH, CELF3 shRNA Plasmid (m): sc-142261-SH, CELF3 shRNA (h) Lentiviral Particles: sc-88405-V and CELF3 shRNA (m) Lentiviral Particles: sc-142261-V.

Molecular Weight of CELF3: 50 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.