

A1BG (C-13): sc-132610

BACKGROUND

A1BG (α 1B-glycoprotein), also known as A1B, ABG, GAB or HYST2477, is a 495 amino acid secreted glycoprotein that contains 5 immunoglobulin (Ig)-like V-type domains and belongs to the immunoglobulin superfamily. At an average concentration of 22mg/dl, A1BG is expressed in normal adult plasma and is thought to be involved in the regulation of cell behavior and cell recognition. In plasma, A1BG specifically binds to human CRISP-3, a member of the cysteine-rich secretory protein (CRISP) family comprised of evolutionarily conserved proteins which are believed to play a role in the innate immune system. Through its association with CRISP-3, A1BG is believed to function in protecting the body from the circulation of free CRISP-3, a circumstance with potentially harmful effects.

REFERENCES

- Ishioka, N., Takahashi, N. and Putnam, F.W. 1986. Amino acid sequence of human plasma α 1B-glycoprotein: homology to the immunoglobulin supergene family. Proc. Natl. Acad. Sci. USA 83: 2363-2367.
- Gahne, B., Juneja, R.K. and Stratil, A. 1987. Genetic polymorphism of human plasma α 1B-glycoprotein: phenotyping by immunoblotting or by a simple method of 2-D electrophoresis. Hum. Genet. 76: 111-115.
- Juneja, R.K., Weitkamp, L.R., Stratil, A., Gahne, B. and Guttormsen, S.A. 1988. Further studies of the plasma α 1B-glycoprotein polymorphism: two new alleles and allele frequencies in Caucasians and in American blacks. Hum. Hered. 38: 267-272.
- Eiberg, H., Bisgaard, M.L. and Mohr, J. 1989. Linkage between α 1B-glycoprotein (A1BG) and Lutheran (LU) red blood group system: assignment to chromosome 19: new genetic variants of A1BG. Clin. Genet. 36: 415-418.
- Juneja, R.K., Saha, N., Tay, J.S., Low, P.S. and Gahne, B. 1994. Distribution of plasma α 1B-glycoprotein (A1BG) polymorphism in several populations of the Indian subcontinent. Ann. Hum. Biol. 21: 443-448.
- Hashim, O.H., Ahmad, F. and Shuib, A.S. 2001. The application of *Artocarpus integer* seed lectin-M in the detection and isolation of selective human serum acute-phase proteins and immunoglobulins. Immunol. Invest. 30: 131-141.
- Sun, D., Zhao, Y.Y., Dai, S.P., Fang, K., Dong, L.Y. and Ding, Q. 2002. Cloning and analysis of human α 1B-glycoprotein precursor gene: a novel member of human immunoglobulin superfamily. Yi Chuan Xue Bao 29: 299-302.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 138670. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Udby, L., Sørensen, O.E., Pass, J., Johnsen, A.H., Behrendt, N., Borregaard, N. and Kjeldsen, L. 2004. Cysteine-rich secretory protein 3 is a ligand of α 1B-glycoprotein in human plasma. Biochemistry 43: 12877-12886.

CHROMOSOMAL LOCATION

Genetic locus: A1BG (human) mapping to 19q13.43.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

A1BG (C-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of A1BG of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-132610 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

A1BG (C-13) is recommended for detection of A1BG of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for A1BG siRNA (h): sc-97518, A1BG shRNA Plasmid (h): sc-97518-SH and A1BG shRNA (h) Lentiviral Particles: sc-97518-V.

Molecular Weight of A1BG: 54 kDa.

Molecular Weight of deglycosylated A1BG: 68 kDa.

Molecular Weight of glycosylated A1BG: 74-80 kDa.

Positive Controls: HeLa nuclear extract: sc-2120, Hep G2 cell lysate: sc-2227 or MCF7 nuclear extract: sc-2149.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfation
Guaranteed

Try **A1BG (F-9): sc-374415** or **A1BG (51A6): sc-135661**, our highly recommended monoclonal alternatives to A1BG (C-13).