G6f (P-13): sc-132697



The Power to Question

BACKGROUND

G6f, also known as lymphocyte antigen 6 complex locus protein G6f, is a single-pass type I membrane protein belonging to the immunoglobulin (Ig) superfamily. Located in the class III region of the major histocompatibility complex (MHC), the G6f gene lies in a cluster of genes encoding cell-surface proteins that play a role in the immune system and cellular recognition. G6f functions as a downstream effector of GRB2 and GRB7, and, in humans, it interacts with GRB2 and GRB7 through the phosphorylation of a tyrosine residue (Tyr 281) in the intracellular tail of G6f. This interaction is also mediated by the SH2 domain of GRB2 and possibly that of GRB7. G6f is a 297 amino acid protein, and it forms a disulfide-linked homodimer.

REFERENCES

- Mungall, A.J., Palmer, S.A., Sims, S.K., Edwards, C.A., Ashurst, J.L., Wilming, L., Jones, M.C., Horton, R., Hunt, S.E., Scott, C.E., Gilbert, J.G.R., Clamp, M.E., Bethel, G., Milne, S., Ainscough, R., Almeida, J.P., Ambrose, K.D., et al. 2003. The DNA sequence and analysis of human chromosome 6. Nature 425: 805-811.
- Vuoristo, M.M., Pappas, J.G., Jansen, V. and Ala-Kokko, L. 2004. A stop codon mutation in COL11A2 induces exon skipping and leads to non-ocular Stickler syndrome. Am. J. Med. Genet. 130A: 160-164.
- McQueen, M.B., Devlin, B., Faraone, S.V., Nimgaonkar, V.L., Sklar, P., Smoller, J.W., Abou Jamra, R., Albus, M., et al. 2005. Combined analysis from eleven linkage studies of bipolar disorder provides strong evidence of susceptibility loci on chromosomes 6q and 8q. Am. J. Hum. Genet. 77: 582-595.
- 4. Batts, K.P. 2007. Iron overload syndromes and the liver. Mod. Pathol. 20 Suppl. 1: S31-S39.
- 5. Olsson, K.S., Ritter, B. and Hansson, N. 2007. The HLA-A1-B8 haplotype hitchhiking with the hemochromatosis mutation: does it affect the phenotype? Eur. J. Haematol. 79: 429-434.

CHROMOSOMAL LOCATION

Genetic locus: LY6G6F (human) mapping to 6p21.33.

SOURCE

G6f (P-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of G6f of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-132697 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

G6f (P-13) is recommended for detection of G6f of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with G6b.

Suitable for use as control antibody for G6f siRNA (h): sc-95654, G6f shRNA Plasmid (h): sc-95654-SH and G6f shRNA (h) Lentiviral Particles: sc-95654-V.


Molecular Weight of G6f: 29 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



G6f (P-13): sc-132697. Western blot analysis of G6f expression in rat small intestine tissue extract ($\bf A$) and Hep G2 whole cell lysate ($\bf B$).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**