

HATH-6 (C-12): sc-132706

BACKGROUND

HATH-6, also known as atonal homolog 8 (ATO8), is a 321 amino acid putative transcription factor. Localized to the nucleus, HATH-6 is expressed in kidney, lung, liver, pancreas, heart and the endothelium of umbilical vessels. HATH-6 is thought to participate in podocyte development in the kidney and may be involved in specification and differentiation of brain neuronal cell lineages. As a transcription factor, HATH-6 contains one basic helix-loop-helix (bHLH) domain. In order to efficiently bind to DNA, HATH-6 must be bound to another bHLH domain containing protein. HATH-6 exists as two isoforms produced by alternative splicing.

REFERENCES

1. Murre, C., McCaw, P.S. and Baltimore, D. 1989. A new DNA binding and dimerization motif in immunoglobulin enhancer binding, daughterless, MyoD, and Myc proteins. *Cell* 56: 777-783.
2. Kato, G.J. and Dang, C.V. 1992. Function of the c-Myc oncoprotein. *FASEB J.* 6: 3065-3072.
3. Inoue, C., Bae, S.K., Takatsuka, K., Inoue, T., Bessho, Y. and Kageyama, R. 2001. Math6, a bHLH gene expressed in the developing nervous system, regulates neuronal versus glial differentiation. *Genes Cells* 6: 977-986.
4. Ledent, V. and Vervoort, M. 2001. The basic helix-loop-helix protein family: comparative genomics and phylogenetic analysis. *Genome Res.* 11: 754-770.
5. Ledent, V., Paquet, O. and Vervoort, M. 2002. Phylogenetic analysis of the human basic helix-loop-helix proteins. *Genome Biol.* 3: RESEARCH0030.
6. Wasserman, S.M., Mehraban, F., Komuves, L.G., Yang, R.B., Tomlinson, J.E., Zhang, Y., Spriggs, F. and Topper, J.N. 2002. Gene expression profile of human endothelial cells exposed to sustained fluid shear stress. *Physiol. Genomics* 12: 13-23.

CHROMOSOMAL LOCATION

Genetic locus: ATO8 (human) mapping to 2p11.2; Atoh8 (mouse) mapping to 6 C1.

SOURCE

HATH-6 (C-12) is available as either goat (sc-132706) or rabbit (sc-132706-R) polyclonal affinity purified antibody raised against a peptide mapping at the C-terminus of HATH-6 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-132706 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-132706 X, 200 µg/0.1 ml.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

HATH-6 (C-12) is recommended for detection of HATH-6 isoforms 1 and 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with HATH-1.

HATH-6 (C-12) is also recommended for detection of HATH-6 isoforms 1 and 2 in additional species, including canine.

Suitable for use as control antibody for HATH-6 siRNA (h): sc-94577, HATH-6 siRNA (m): sc-145899, HATH-6 shRNA Plasmid (h): sc-94577-SH, HATH-6 shRNA Plasmid (m): sc-145899-SH, HATH-6 shRNA (h) Lentiviral Particles: sc-94577-V and HATH-6 shRNA (m) Lentiviral Particles: sc-145899-V.

HATH-6 (C-12) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of HATH-6: 35 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227, K-562 whole cell lysate: sc-2203 or NIH/3T3 whole cell lysate: sc-2210.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



HATH-6 (C-12): sc-132706. Western blot analysis of HATH-6 expression in K-562 (A), Hep G2 (B) and NIH/3T3 (C) whole cell lysates and mouse brain (D), human liver (E) and rat pancreas (F) tissue extracts.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.