

RNF135 (P-13): sc-132800

BACKGROUND

The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain this conserved domain are generally involved in the ubiquitination pathway of protein degradation. RNF135 (RING finger protein 135), also known as L13, is a 432 amino acid protein that contains one RING-type zinc finger and one SPRY domain. Via its RING-type zinc finger, RNF135 may play a role in transcriptional regulation and protein degradation events. Defects in the gene encoding RNF135 are the cause of RNF135-related overgrowth syndrome which is characterized by learning disabilities, facial dysmorphism and increased weight and height. Multiple isoforms of RNF135 exist due to alternative splicing events.

REFERENCES

- Borden, K.L. and Freemont, P.S. 1996. The RING finger domain: a recent example of a sequence-structure family. *Curr. Opin. Struct. Biol.* 6: 395-401.
- Lorick, K.L., Jensen, J.P., Fang, S., Ong, A.M., Hatakeyama, S. and Weissman, A.M. 1999. RING fingers mediate ubiquitin-conjugating enzyme (E2)-dependent ubiquitination. *Proc. Natl. Acad. Sci. USA* 96: 11364-11369.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 611358. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Jenne, D.E., Tinschert, S., Dorschner, M.O., Hameister, H., Stephens, K. and Kehrer-Sawatzki, H. 2003. Complete physical map and gene content of the human NF-1 tumor suppressor region in human and mouse. *Genes Chromosomes Cancer* 37: 111-120.
- Colland, F., Jacq, X., Trouplin, V., Mouglin, C., Groizeleau, C., Hamburger, A., Meil, A., Wojcik, J., Legrain, P. and Gauthier, J.M. 2004. Functional proteomics mapping of a human signaling pathway. *Genome Res.* 14: 1324-1332.
- Penengo, L., Mapelli, M., Murachelli, A.G., Confalonieri, S., Magri, L., Musacchio, A., Di Fiore, P.P., Polo, S. and Schneider, T.R. 2006. Crystal structure of the ubiquitin binding domains of Rabex-5 reveals two modes of interaction with ubiquitin. *Cell* 124: 1183-1195.
- Douglas, J., Cilliers, D., Coleman, K., Tatton-Brown, K., Barker, K., Bernhard, B., Burn, J., Huson, S., Josifova, D., Lacombe, D., Malik, M., Mansour, S., et al. 2007. Mutations in RNF135, a gene within the NF-1 microdeletion region, cause phenotypic abnormalities including overgrowth. *Nat. Genet.* 39: 963-965.

CHROMOSOMAL LOCATION

Genetic locus: RNF135 (human) mapping to 17q11.2.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

SOURCE

RNF135 (P-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of RNF135 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-132800 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

RNF135 (P-13) is recommended for detection of RNF135 isoforms 1 and 2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other RNF family members.

Suitable for use as control antibody for RNF135 siRNA (h): sc-94030, RNF135 shRNA Plasmid (h): sc-94030-SH and RNF135 shRNA (h) Lentiviral Particles: sc-94030-V.

Molecular Weight of RNF135: 48 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.