

# Chfr (N-15): sc-13287

## BACKGROUND

The forkhead-associated (FHA) domain was initially identified in transcription factors that have forkhead DNA-binding domains and in protein kinases, but many cell-cycle checkpoint proteins, including chfr (checkpoint with forkhead and ring finger domains) contain FHA domains. Chfr defines a checkpoint that delays entry into metaphase in response to mitotic stress. Normal primary cells and tumor cell lines that express wild-type chfr exhibit delayed entry into metaphase when centrosome separation is inhibited by mitotic stress. Additionally, chfr seems to be required for delaying prophase in human cells. The sequence of chfr is similar to that of the fission yeast DMA1, which is involved in a later mitotic checkpoint that delays a cell's exit from mitosis in response to spindle damage.

## REFERENCES

- Jha, M.N., et al. 1994. Cell cycle arrest by Colcemid differs in human normal and tumor cells. *Cancer Res.* 54: 5011-5015.
- Hofmann, K. and Bucher, P. 1995. The FHA domain: a putative nuclear signalling domain found in protein kinases and transcription factors. *Trends Biochem. Sci.* 20: 347-349.
- Murone, M. and Simanis, V. 1996. The fission yeast dma1 gene is a component of the spindle assembly checkpoint, required to prevent septum formation and premature exit from mitosis if spindle function is compromised. *EMBO J.* 15: 6605-6616.
- Cortez, D. and Elledge, S.J. 2000. Conducting the mitotic symphony. *Nature* 406: 354-356.
- Scolnick, D.M. and Halazonetis, T.D. 2000. Chfr defines a mitotic stress checkpoint that delays entry into metaphase. *Nature* 406: 430-435.
- Tokunaga, E., et al. 2006. Aberrant hypermethylation of the promoter region of the CHFR gene is rare in primary breast cancer. *Breast Cancer Res. Treat.* 97: 199-203.
- Kobayashi, C., et al. 2006. Aberrant expression of CHFR in malignant peripheral nerve sheath tumors. *Mod. Pathol.* 19: 524-532.
- Koga, Y., et al. 2006. The significance of aberrant CHFR methylation for clinical response to microtubule inhibitors in gastric cancer. *J. Gastroenterol.* 41: 133-139.
- Yanokura, M., et al. 2007. Relationship of aberrant DNA hypermethylation of CHFR with sensitivity to taxanes in endometrial cancer. *Oncol. Rep.* 17: 41-48.

## CHROMOSOMAL LOCATION

Genetic locus: CHFR (human) mapping to 12q24.33.

## SOURCE

Chfr (N-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Chfr of human origin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-13287 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

Chfr (N-15) is recommended for detection of Chfr of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Chfr (N-15) is also recommended for detection of Chfr in additional species, including equine, bovine, porcine and avian.

Suitable for use as control antibody for Chfr siRNA (h): sc-37567, Chfr shRNA Plasmid (h): sc-37567-SH and Chfr shRNA (h) Lentiviral Particles: sc-37567-V.


Molecular Weight of Chfr isoforms: 73/72/69/64 kDa.

Positive Controls: ES-2 cell lysate: sc-24674.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

## DATA



Chfr (N-15): sc-13287. Immunoperoxidase staining of formalin fixed, paraffin-embedded human bone marrow tissue showing nuclear and cytoplasmic staining of hematopoietic cells.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\* Stable for one year from the date of shipment. Non-hazardous. No MSDS required.