

ZNF213 (Q-13): sc-132912

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF213 (zinc finger protein 213), also known as CR53 or ZKSCAN21, is a 459 amino acid protein that contains 5 C₂H₂-type zinc fingers, one KRAB domain and one SCAN box domain. Localized to the nucleus and expressed at high levels in the testis, ZNF213 belongs to the krüppel C₂H₂-type zinc-finger protein family and is thought to be involved in transcriptional regulation.

REFERENCES

- Bernot, A., Heilig, R., Clepet, C., Smaoui, N., Da Silva, C., Petit, J.L., Devaud, C., Chiannikulchai, N., Fizames, C., Samson, D., Cruaud, C., Caloustian, C., Gyapay, G., Delpech, M. and Weissenbach, J. 1998. A transcriptional Map of the FMF region. *Genomics* 50: 147-160.
- Chen, X., Hamon, M., Deng, Z., Centola, M., Sood, R., Taylor, K., Kastner, D.L. and Fischel-Ghodsian, N. 1999. Identification and characterization of a zinc finger gene (ZNF213) from 16p13.3. *Biochim. Biophys. Acta* 1444: 218-230.
- Rousseau-Merck, M.F., Koczan, D., Legrand, I., Möller, S., Autran, S. and Thiesen, H.J. 2002. The KOX zinc finger genes: genome wide mapping of 368 ZNF PAC clones with zinc finger gene clusters predominantly in 23 chromosomal loci are confirmed by human sequences annotated in Ensembl. *Cytogenet. Genome Res.* 98: 147-153.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 608387. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Nakamura, M., Runko, A.P. and Sagerström, C.G. 2004. A novel subfamily of zinc finger genes involved in embryonic development. *J. Cell. Biochem.* 93: 887-895.
- Englbrecht, C.C., Schoof, H. and Böhm, S. 2004. Conservation, diversification and expansion of C₂H₂ zinc finger proteins in the *Arabidopsis thaliana* genome. *BMC Genomics* 5: 39-39.

CHROMOSOMAL LOCATION

Genetic locus: ZNF213 (human) mapping to 16p13.3.

SOURCE

ZNF213 (Q-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ZNF213 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-132912 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ZNF213 (Q-13) is recommended for detection of ZNF213 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other ZNF family members.

Suitable for use as control antibody for ZNF213 siRNA (h): sc-93233, ZNF213 shRNA Plasmid (h): sc-93233-SH and ZNF213 shRNA (h) Lentiviral Particles: sc-93233-V.


Molecular Weight of ZNF213: 51 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201 or Hep G2 cell lysate: sc-2227.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



ZNF213 (Q-13): sc-132912. Immunoperoxidase staining of formalin fixed, paraffin-embedded human nasopharynx tissue showing nuclear staining of respiratory epithelial cells.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.