


γ PAK (F-12): sc-133113



The Power to Question

BACKGROUND

Three isoforms of serine/threonine kinases, designated α PAK p68, β PAK p65 and γ PAK p62, have been shown to exhibit a high degree of sequence homology with the *S. cerevisiae* kinase Ste 20, involved in pheromone signaling. The α , β and γ PAK isoforms complex specifically with Rac1 and Cdc42 in their active GTP-bound state, inhibiting their intrinsic GTPase activity leading to their autophosphorylation. There are eight sites of autophosphorylation on γ PAK, including Ser 19, Ser 141 and Thr 402, and phosphorylation of Ser 141 and Thr 402 is correlated with γ PAK activation. Once phosphorylated and their affinity for Rac/Cdc42 reduced, the PAK isoforms disassociate from the complex to seek downstream substrates. One such putative substrate is Mek kinase, an upstream effector of Mek4 which is involved in the JNK signaling pathway. While the PAK isoforms interact in a GTP-dependent manner with Rac1 and Cdc42, they do not interact with Rho.

REFERENCES

- Didsbury, J., et al. 1989. Rac, a novel Ras-related family of proteins that are botulinum toxic substrates. *J. Biol. Chem.* 264: 16378-16382.
- Shinjo, K., et al. 1990. Molecular cloning of the gene for the human placental GTP-binding protein G_p (G25K): identification of this GTP-binding protein as the human homolog of the yeast cell-division-cycle protein CDC42. *Proc. Natl. Acad. Sci. USA* 87: 9853-9857.
- Boguski, M.S. and McCormick, F. 1993. Proteins regulating Ras and its relatives. *Nature* 366: 643-654.
- Manser, E., et al. 1994. A brain serine/threonine protein kinase activated by Cdc42 and Rac1. *Nature* 367: 40-46.
- Yan, M., et al. 1994. Activation of stress-activated protein kinase by MEKK1 phosphorylation of its activator SEK1. *Nature* 372: 798-800.
- Minden, A., et al. 1994. Differential activation of ERK and JNK mitogen-activated protein kinases by Raf-1 and MEKK. *Science* 266: 1719-1723.
- Martin, G.A., et al. 1995. A novel serine kinase activated by rac1/CDC42Hs-dependent autophosphorylation is related to PAK65 and STE20. *EMBO J.* 14: 1970-1978.

CHROMOSOMAL LOCATION

Genetic locus: PAK2 (human) mapping to 3q29; Pak2 (mouse) mapping to 16 B2.

SOURCE

γ PAK (F-12) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 2-41 at the N-terminus of γ PAK of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-133113 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

γ PAK (F-12) is recommended for detection of γ PAK p62 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

γ PAK (F-12) is also recommended for detection of γ PAK p62 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for γ PAK siRNA (h): sc-36183, γ PAK siRNA (m): sc-36184, γ PAK shRNA Plasmid (h): sc-36183-SH, γ PAK shRNA Plasmid (m): sc-36184-SH, γ PAK shRNA (h) Lentiviral Particles: sc-36183-V and γ PAK shRNA (m) Lentiviral Particles: sc-36184-V.


Molecular Weight of γ PAK: 62 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201, JAR cell lysate: sc-2276 or Jurkat whole cell lysate: sc-2204.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



γ PAK (F-12): sc-133113. Western blot analysis of γ PAK expression in JAR (A) and Jurkat (B) whole cell lysates.

SELECT PRODUCT CITATIONS

- Kathania, M., et al. 2022. Pak2-mediated phosphorylation promotes ROR γ t ubiquitination and inhibits colonic inflammation. *Cell Rep.* 40: 111345.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.