SANTA CRUZ BIOTECHNOLOGY, INC.

UKHC (F-9): sc-133185



BACKGROUND

The kinesin motor proteins include at least two forms of conventional kinesin encoded by different genes and designated as ubiquitous kinesin, which is expressed in all cells and tissues, or neuronal kinesin, which is expressed exclusively in neural cells. Kinesin is a microtubule associated protein comprised of three different structural domains. A considerable globular N-terminal domain regulates the hydrolysis of ATP and also microtubule binding while central coiled-coil domains promote heavy chain dimerization. Lastly, small globular C-terminal domains interact with kinesin light chains, membranous organelles and vesicles. Expression of ubiquitous kinesin heavy chain, also designated UKHC, is found subcellularly in areas of heavy vesicular trafficking such as the microtubule pathways of neural cells and also the Golgi of non-neural cell types.

REFERENCES

- Ingold, A.L., et al. 1989. Inhibition of kinesin-driven microtubule motility by monoclonal antibodies to kinesin heavy chains. J. Cell Biol. 107: 2657-2667.
- Wright, B.D., et al. 1991. Subcellular localization and sequence of sea urchin kinesin heavy chain: evidence for its association with membranes in the mitotic apparatus and interphase cytoplasm. J. Cell Biol. 113: 817-833.
- 3. Marks, D.L., et al. 1995. Association of kinesin with the Golgi apparatus in rat hepatocytes. J. Cell Sci. 107: 2417-2426.
- Bi, G.Q., et al. 1997. Kinesin- and myosin-driven steps of vesicle recruitment for Ca²⁺-regulated exocytosis. J. Cell Biol. 138: 999-1008.
- DeLuca, J.G., et al. 2001. Purification and characterization of native conventional kinesin, HSET, and CENP-E from mitotic HeLa cells. J. Biol. Chem. 276: 28014-28021.
- Fichera, M., et al. 2004. Evidence of kinesin heavy chain (KIF5A) involvement in pure hereditary spastic paraplegia. Neurology 63: 1108-1110.
- Li, C., et al. 2006. A family-based association study of kinesin heavy chain member 2 gene (KIF2) and schizophrenia. Neurosci. Lett. 407: 151-155.
- Glater, E.E., et al. 2006. Axonal transport of mitochondria requires milton to recruit kinesin heavy chain and is light chain independent. J. Cell Biol. 173: 545-557.

CHROMOSOMAL LOCATION

Genetic locus: KIF5B (human) mapping to 10p11.22; Kif5b (mouse) mapping to 18 A1.

SOURCE

UKHC (F-9) is a mouse monoclonal antibody raised against amino acids 691-740 mapping within an internal region of UKHC of human origin.

PRODUCT

Each vial contains 200 μg IgG_1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

UKHC (F-9) is recommended for detection of ubiquitous kinesin heavy chain (UKHC) of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for UKHC siRNA (h): sc-36777, UKHC siRNA (m): sc-36778, UKHC shRNA Plasmid (h): sc-36777-SH, UKHC shRNA Plasmid (m): sc-36778-SH, UKHC shRNA (h) Lentiviral Particles: sc-36777-V and UKHC shRNA (m) Lentiviral Particles: sc-36778-V.

Molecular Weight of UKHC: 120 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Hep G2 cell lysate: sc-2227 or Jurkat whole cell lysate: sc-2204.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG K BP-HRP: sc-516102 or m-lgG K BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-lgG K BP-FITC: sc-516140 or m-lgG K BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA





UKHC (F-9): sc-133185. Western blot analysis of UKHC expression in A-431 (**A**), HeLa (**B**), Hep G2 (**C**) and Jurkat (**D**) whole cell lysates.

UKHC (F-9): sc-133185. Western blot analysis of UKHC expression in Jurkat (**A**), NCI-H1299 (**B**) and Caco-2 (**C**) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.