

Espin (C-12): sc-133322

BACKGROUND

Espin (ESPN), also known as autosomal recessive deafness type 36 protein (DFNB36), is an 854 amino acid cytoplasmic protein that contains nine ANK repeats and one WH2 domain. The WH2 domain of Espin binds Actin monomers and mediates the assembly of the Actin bundle. This interaction plays a major role in the moderation of the organization, dynamics and signaling capacities of the Actin filament-rich specializations that regulate sensory transduction in various sensory cells. Defects in Espin are the cause of non-syndromic sensorineural deafness autosomal recessive type 36 (DFNB36), a sensorineural hearing loss caused by damage to the neural receptors of the inner ear, the nerve pathways to the brain or the region of the brain responsible for sound. In humans, Espin is expressed as two isoforms produced by alternative splicing and has been found to interact with IRSp53 and Profilin-2. In rodents, four major isoforms ranging from approximately 110 to 25 kDa have been identified with additional splice variants possible.

REFERENCES

- Bartles, J.R., Wierda, A. and Zheng, L. 1996. Identification and characterization of Espin, an Actin-binding protein localized to the F-Actin-rich junctional plaques of Sertoli cell ectoplasmic specializations. *J. Cell Sci.* 109: 1229-1239.
- Zheng, L., Sekerková, G., Vranich, K., Tilney, L.G., Mugnaini, E. and Bartles, J.R. 2000. The deaf jerker mouse has a mutation in the gene encoding the Espin Actin-bundling proteins of hair cell stereocilia and lacks Espins. *Cell* 102: 377-385.7
- Naz, S., Griffith, A.J., Riazuddin, S., Hampton, L.L., Battey, J.F., Khan, S.N., Riazuddin, S., Wilcox, E.R. and Friedman, T.B. 2004. Mutations of ESPN cause autosomal recessive deafness and vestibular dysfunction. *J. Med. Genet.* 41: 591-595.
- Loomis, P.A., Kelly, A.E., Zheng, L., Changyaleket, B., Sekerková, G., Mugnaini, E., Ferreira, A., Mullins, R.D. and Bartles, J.R. 2006. Targeted wild-type and jerker Espins reveal a novel, WH2-domain-dependent way to make Actin bundles in cells. *J. Cell Sci.* 119: 1655-1665.
- Donaudy, F., Zheng, L., Ficarella, R., Ballana, E., Carella, M., Melchionda, S., Estivill, X., Bartles, J.R. and Gasparini, P. 2006. Espin gene (ESPN) mutations associated with autosomal dominant hearing loss cause defects in microvillar elongation or organisation. *J. Med. Genet.* 43: 157-161.
- Primiani, N., Gregory, M., Dufresne, J., Smith, C.E., Liu, Y.L., Bartles, J.R., Cyr, D.G. and Hermo, L. 2007. Microvillar size and Espin expression in principal cells of the adult rat epididymis are regulated by androgens. *J. Androl.* 28: 659-669.
- Boulouiz, R., Li, Y., Soualhine, H., Abidi, O., Chafik, A., Nürnberg, G., Becker, C., Nürnberg, P., Kubisch, C., Wollnik, B. and Barakat, A. 2008. A novel mutation in the Espin gene causes autosomal recessive nonsyndromic hearing loss but no apparent vestibular dysfunction in a Moroccan family. *Am. J. Med. Genet. A.* 146A: 3086-3089.

CHROMOSOMAL LOCATION

Genetic locus: ESPN (human) mapping to 1p36.31; Espn (mouse) mapping to 4 E2.

SOURCE

Espin (C-12) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping at the C-terminus of Espin of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-133322 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Espin (C-12) is recommended for detection of Espin of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:50-1:500), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:25, dilution range 1:25-1:250) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Espin (C-12) is also recommended for detection of Espin in additional species, including equine, canine and porcine.


Suitable for use as control antibody for Espin siRNA (h): sc-78697, Espin siRNA (m): sc-40510, Espin shRNA Plasmid (h): sc-78697-SH, Espin shRNA Plasmid (m): sc-40510-SH, Espin shRNA (h) Lentiviral Particles: sc-78697-V and Espin shRNA (m) Lentiviral Particles: sc-40510-V.

Molecular Weight of Espin isoform 1: 110 kDa.

Molecular Weight of Espin mouse variant: 25 kDa.

Positive Controls: Espin (m): 293T Lysate: sc-125310 or rat testis extract: sc-2400.

DATA



Espin (C-12): sc-133322. Western blot analysis of Espin expression in non-transfected: sc-117752 (A) and mouse Espin transfected: sc-125310 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.