



SCLT1 (N-14): sc-133432

BACKGROUND

SCLT1 (sodium channel and clathrin linker 1), also known as CAP1A or hCAP-1A, is a 688 amino acid cytoplasmic protein that acts as a linker between the voltage-gated sodium channel, Na⁺ CP type X α , and clathrin. SCLT1 is abundantly expressed in DRG (dorsal root ganglia) neurons and co-localizes with Na⁺ CP type X α . SCLT1 regulates Na⁺ CP type X α channel activity by promoting channel internalization. SCLT1 exists as four alternatively spliced isoforms and is encoded by a gene located on human chromosome 4, which encodes nearly 6% of the human genome and has the largest gene deserts (regions of the genome with no protein encoding genes) of all human chromosomes. Defects in some of the genes located on chromosome 4 are associated with Huntington's disease, Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.

REFERENCES

1. Lemmon, S.K. 2001. Clathrin uncoating: Auxilin comes to life. *Curr. Biol.* 11: R49-R52.
2. Rougier, J.S., van Bemmelen, M.X., Bruce, M.C., Jespersen, T., Gavillet, B., Apotheloz, F., Cordonier, S., Staub, O., Rotin, D. and Abriel, H. 2005. Molecular determinants of voltage-gated sodium channel regulation by the NEDD4/NEDD4-like proteins. *Am. J. Physiol., Cell Physiol.* 288: C692-C701.
3. Liu, C., Cummins, T.R., Tyrrell, L., Black, J.A., Waxman, S.G. and Dib-Hajj, S.D. 2005. CAP-1A is a novel linker that binds clathrin and the voltage-gated sodium channel Na(v)1.8. *Mol. Cell. Neurosci.* 28: 636-649.
4. Hillier, L.W., Graves, T.A., Fulton, R.S., Fulton, L.A., Pepin, K.H., Minx, P., Wagner-McPherson, C., Layman, D., Wylie, K., Sekhon, M., Becker, M.C., Fewell, G.A., et al. 2005. Generation and annotation of the DNA sequences of human chromosomes 2 and 4. *Nature* 434: 724-731.
5. Cowan, C.M. and Raymond, L.A. 2006. Selective neuronal degeneration in Huntington's disease. *Curr. Top. Dev. Biol.* 75:25-71.
6. Cummins, T.R., Sheets, P.L. and Waxman, S.G. 2007. The roles of sodium channels in nociception: Implications for mechanisms of pain. *Pain* 131: 243-257.
7. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 611399. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: SCLT1 (human) mapping to 4q28.2.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

SOURCE

SCLT1 (N-14) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping near the N-terminus of SCLT1 of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-133432 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SCLT1 (N-14) is recommended for detection of SCLT1 isoforms 1-4 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:50-1:500), immunofluorescence (starting dilution 1:25, dilution range 1:25-1:250) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SCLT1 siRNA (h): sc-89139, SCLT1 shRNA Plasmid (h): sc-89139-SH and SCLT1 shRNA (h) Lentiviral Particles: sc-89139-V.

Molecular Weight of SCLT1: 81 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.