

FIR (E-23): sc-133583

BACKGROUND

Activation of FUSE, the far-upstream element, is required for the proper expression of the mammalian gene c-Myc. The binding of FBP (FUSE-binding protein) to FUSE is necessary for c-Myc expression. The FBP interacting repressor, FIR, binds to the central DNA-binding domain of FBP and can serve as an overriding negative regulator of c-Myc promoter activity. FIR interacts with the TFIID complex, which is a multifunctional, multisubunit RNA polymerase II transcription factor that interacts with several DNA-binding transactivators. FIR blocks activator-dependent, but not basal transcription through TFIID. FIR shares identity with seven in absentia (sia) binding protein 1. FIR is expressed in spleen, thymus, prostate, small intestine, colon, and peripheral blood leukocytes, and with relatively higher levels of expression in testis and ovary.

REFERENCES

1. Avigan, M.I., Stober, B. and Levens, D., 1990. A far upstream element stimulates c-Myc expression in undifferentiated leukemia cells. *J. Biol. Chem.* 265: 18538-18545.
2. Duncan, R.D., Bazar, L., Michelotti, G., Tomonaga, T., Krutzsch, H., Avigan, M. and Levens, D. 1994. A sequence-specific, single strand binding protein activates the far upstream of c-Myc and defines a new DNA binding motif. *Genes Dev.* 8: 465-480.
3. Bazar, L., Harris, V., Sunitha, I., Hartmann, D. and Avigan, M. 1995. A transactivator of c-Myc is coordinately regulated with the proto-oncogene during cellular growth. *Oncogene* 10: 2229-2238.
4. Frit, P., Bergmann, E. and Egly, J-M. 1999. Transcription factor IIIH: a key player in the cellular response to DNA damage. *Biochimie* 81: 27-38.
5. Haile, D.T. and Parvin, J.D. 1999. Activation of transcription *in vitro* by the BRCA1 carboxyl-terminal domain. *J. Biol. Chem.* 274: 2113-2117.
6. Liu, J., Liusheng, H., Collins, I., Ge, H., Libuitti, D., Li, J., Egly, J-M. and Levens, D. 2000. The FBP interacting repressor targets TFIID to inhibit activated transcription. *Mol. Cell* 5: 331-341.

CHROMOSOMAL LOCATION

Genetic locus: PUF60 (human) mapping to 8q24.3; Puf60 (mouse) mapping to 15 D3.

SOURCE

FIR (E-23) is an affinity purified rabbit polyclonal antibody raised against synthetic FIR peptide of human origin.

PRODUCT

Each vial contains 50 µg IgG in 500 µl PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

FIR (E-23) is recommended for detection of FIR of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FIR siRNA (h): sc-105353, FIR siRNA (m): sc-145186, FIR shRNA Plasmid (h): sc-105353-SH, FIR shRNA Plasmid (m): sc-145186-SH, FIR shRNA (h) Lentiviral Particles: sc-105353-V and FIR shRNA (m) Lentiviral Particles: sc-145186-V.


Molecular Weight of FIR: 60 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



FIR (E-23): sc-133583. Western blot analysis of FIR expression in HeLa whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **FIR (E-6): sc-398799** or **FIR (B-5): sc-398785**, our highly recommended monoclonal alternatives to FIR (E-23).