

FO XK2 (G-23): sc-133593

BACKGROUND

The forkhead-box (FOX) genes comprise a superfamily of at least 43 members that encode proteins which are involved in transcriptional regulation and may be associated with the pathogenesis of various cancers. FOXK2 (forkhead box K2), also known as ILF or ILF1, is a 660 amino acid protein that localizes to the nucleus and contains one FHA domain and one forkhead DNA-binding domain. Expressed in lymphoid and non-lymphoid cells, FOXK2 uses magnesium as a cofactor and binds to purine-rich motifs in select promoters, possibly playing a role in the regulation of viral and cellular promoter elements. FOXK2 exists as multiple alternatively spliced isoforms and is subject to DNA damage-dependent phosphorylation, probably by Atm or ATR.

REFERENCES

- Durand, D.B., Shaw, J.P., Bush, M.R., Replogle, R.E., Belagaje, R. and Crabtree, G.R. 1988. Characterization of antigen receptor response elements within the interleukin-2 enhancer. *Mol. Cell. Biol.* 8: 1715-1724.
- Li, C., Lai, C.F., Sigman, D.S. and Gaynor, R.B. 1991. Cloning of a cellular factor, interleukin binding factor, that binds to NFAT-like motifs in the human immunodeficiency virus long terminal repeat. *Proc. Natl. Acad. Sci. USA* 88: 7739-7743.
- Li, C., Lusic, A.J., Sparkes, R., Nirula, A. and Gaynor, R. 1992. Characterization and chromosomal mapping of the gene encoding the cellular DNA binding protein ILF. *Genomics* 13: 665-671.
- Nirula, A., Moore, D.J. and Gaynor, R.B. 1997. Constitutive binding of the transcription factor interleukin-2 (IL-2) enhancer binding factor to the IL-2 promoter. *J. Biol. Chem.* 272: 7736-7745.
- Liu, P.P., Chen, Y.C., Li, C., Hsieh, Y.H., Chen, S.W., Chen, S.H., Jeng, W.Y. and Chuang, W.J. 2002. Solution structure of the DNA-binding domain of interleukin enhancer binding factor 1 (FOXK1a). *Proteins* 49: 543-553.
- Katoh, M. and Katoh, M. 2004. Human FOX gene family. *Int. J. Oncol.* 25: 1495-1500.
- Tsai, K.L., Huang, C.Y., Chang, C.H., Sun, Y.J., Chuang, W.J. and Hsiao, C.D. 2006. Crystal structure of the human FOXK1a-DNA complex and its implications on the diverse binding specificity of winged helix/forkhead proteins. *J. Biol. Chem.* 281: 17400-17409.
- Online Mendelian Inheritance in Man, OMIM™. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 147685. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: FOXK2 (human) mapping to 17q25.3.

SOURCE

FOXK2 (G-23) is an affinity purified rabbit polyclonal antibody raised against synthetic FOXK2 peptide of human origin.

PRODUCT

Each vial contains 50 µg IgG in 500 µl PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

APPLICATIONS

FOXK2 (G-23) is recommended for detection of FOXK2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FOXK2 siRNA (h): sc-94054, FOXK2 shRNA Plasmid (h): sc-94054-SH and FOXK2 shRNA (h) Lentiviral Particles: sc-94054-V.


Molecular Weight of FOXK2: 69 kDa.

Positive Controls: Hep G2 Cell Lysate: sc-2227.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



FOXK2 (G-23): sc-133593. Western blot analysis of FOXK2 expression in Hep G2 whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.