

Gcom1 (L-22): sc-133612

BACKGROUND

Glutamate receptors mediate most excitatory neurotransmission in the brain and play an important role in neural plasticity, neural development and neurodegeneration. Ionotropic glutamate receptors are categorized into NMDA receptors and kainate/AMPA receptors, both of which contain glutamate-gated, cation-specific ion channels. Synaptic and extrasynaptic NMDA receptors have been shown to have opposite effects on neuronal survival, CREB function and gene regulation. Gcom1 (GRINL1A complex locus protein 1), also known as GUP (GRINL1A upstream protein) and Gcom (GRINL1A combined protein), is a 466 amino acid protein that is a component of the GRINL1A complex transcription unit, which is thought to be involved in the modulation of glutamatergic neurotransmission through interaction with the NR1 subunit of the NMDA receptor. Gcom1 is expressed in small intestine, lung, liver, heart, skeletal muscle, testis and prostate and also colocalizes with NR1 in cortical and hippocampal neurons. There are eleven isoforms of Gcom1 that are produced as a result of alternative splicing events.

REFERENCES

- Roginski, R.S., Mohan Raj, B.K., Finkernagel, S.W. and Sciorra, L.J. 2001. Assignment of an ionotropic glutamate receptor-like gene (GRINL1A) to human chromosome 15q22.1 by *in situ* hybridization. *Cytogenet. Cell Genet.* 93: 143-144.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606485. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- McIlhinney, R.A., Philipps, E., Le Bourdelles, B., Grimwood, S., Wafford, K., Sandhu, S. and Whiting, P. 2003. Assembly of N-methyl-D-aspartate (NMDA) receptors. *Biochem. Soc. Trans.* 31: 865-868.
- Black, D.L. and Grabowski, P.J. 2003. Alternative pre-mRNA splicing and neuronal function. *Prog. Mol. Subcell. Biol.* 31: 187-216.
- Roginski, R.S., Mohan Raj, B.K., Birditt, B. and Rowen, L. 2004. The human GRINL1A gene defines a complex transcription unit, an unusual form of gene organization in eukaryotes. *Genomics* 84: 265-276.
- Vazhappilly, R. and Sucher, N.J. 2004. Translational regulation of the N-methyl-D-aspartate receptor subunit NR1. *Neurosignals* 13: 190-193.
- Gibb, A.J. 2004. NMDA receptor subunit gating—uncovered. *Trends Neurosci.* 27: 7-10.

CHROMOSOMAL LOCATION

Genetic locus: Gcom1 (human) mapping to 15q21.3.

SOURCE

Gcom1 (L-22) is an affinity purified rabbit polyclonal antibody raised against synthetic Gcom1 peptide of human origin.

PRODUCT

Each vial contains 50 µg IgG in 500 µl PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

APPLICATIONS

Gcom1 (L-22) is recommended for detection of Gcom1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); not reactive with GRINL1A complex locus downstream isoforms, including Gdown1, Gdown3, Gdown4, and Gdown6.

Suitable for use as control antibody for Gcom1 siRNA (h): sc-90054, Gcom1 shRNA Plasmid (h): sc-90054-SH and Gcom1 shRNA (h) Lentiviral Particles: sc-90054-V.


Molecular Weight of Gcom1: 54 kDa.

Positive Controls: 721 B whole cell lysate.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



Gcom1 (L-22): sc-133612. Western blot analysis of Gcom1 expression in 721 B whole cell lysate.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.