

Glut10 (Q-22): sc-133627

BACKGROUND

Glucose serves as the major energy substrate of mammalian cells and is fundamental to metabolism. Glucose passage across cell membranes is mediated by a family of transporters termed glucose transporters, or Gluts, which are characterized by the presence of 12 membrane-spanning helices. The Glut family is divided into three subfamilies: class I (previously known as glucose transporters), which includes Glut1, Glut2, Glut3 and Glut4; class II (previously known as fructose transporters), which includes Glut5, Glut7, Glut9 and Glut11; and class III, which includes Glut6, Glut8, Glut10, Glut12 and the myo-inositol transporter HMIT1. Glut10 is a 541 amino acid facilitative glucose transporter expressed in high amounts in liver and pancreas. It contains 12 transmembrane domains, with a hydrophilic intracellular loop between helices 6 and 7, and a potential N-linked glycosylation site with a large extracellular loop between helices 9 and 10. The gene for Glut 10, SLC2A10, maps to chromosome 20q13.1, a region that is linked to type 2 diabetes.

REFERENCES

- Shikhman, A., Brinson, D., Valbracht, J. and Lotz, M. 2001. Cytokine regulation of facilitated glucose transport in human articular chondrocytes. *J. Immunology* 167: 7001-7008.
- Joost, H. and Thorens, B. 2001. The extended Glut-family of sugar/polyol transport facilitators: nomenclature, sequence characteristics and function of its novel members. *Mol. Membr. Biol.* 18: 247-256.
- McVie-Wylie, A.J., Lamson, D.R. and Chen Y.T. 2001. Molecular cloning of a novel member of the Glut family of transporters, SLC2A10 (Glut10), localized on chromosome 20q13.1: a candidate gene for NIDDM susceptibility. *Genomics* 72: 113-117.
- Dawson, P.A., Mychaleckyj, J.C., Fossey, S.C., Mihic, S.J., Craddock, A.L. and Bowden, D.W. 2001. Sequence and functional analysis of Glut10: a glucose transporter in the type 2 diabetes-linked region of chromosome 20q12-13.1. *Mol. Genet. Metab.* 74: 186-199.
- Andersen, G., Rose, C.S., Hamid, Y.H., Drivsholm, T., Borch-Johnsen, K., Hansen, T. and Pedersen, O. 2003. Genetic variation of the Glut10 glucose transporter (SLC2A10) and relationships to type 2 diabetes and intermediary traits. *Diabetes* 52: 2445-2448.

CHROMOSOMAL LOCATION

Genetic locus: SLC2A10 (human) mapping to 20q13.12.

SOURCE

Glut10 (Q-22) is a Protein A purified rabbit polyclonal antibody raised against synthetic Glut10 peptide of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Glut10 (Q-22) is recommended for detection of Glut10 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Glut10 siRNA (h): sc-60697, Glut10 shRNA Plasmid (h): sc-60697-SH and Glut10 shRNA (h) Lentiviral Particles: sc-60697-V.


Molecular Weight of Glut10: 57 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



Glut10 (Q-22): sc-133627. Western blot analysis of Glut10 expression in Jurkat whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Glut10 (H-10): sc-398495**, our highly recommended monoclonal alternative to Glut10 (Q-22).