

HFE (W-24): sc-133654

BACKGROUND

The features of hemochromatosis include cirrhosis of the liver, diabetes, hypermelanotic pigmentation of the skin, and heart failure. Since hemochromatosis is a relatively easily treated disorder if diagnosed, this is a form of preventable cancer. The HFE protein, which is defective in hereditary hemochromatosis, normally is expressed in crypt enterocytes of the duodenum where it has a unique, predominantly intracellular localization. In placenta, the HFE protein co-localizes with and forms a stable association with the transferrin receptor (TFR), providing a link between the HFE protein and iron transport. Immunocytochemistry shows that the HFE protein and TFR both are expressed in the crypt enterocytes. Western blots show that, as is the case in human placenta, the HFE protein in crypt enterocytes is physically associated with the TFR and with β 2-microglobulin. It is proposed that HFE has two mutually exclusive activities in cells: inhibition of uptake or inhibition of release of iron and that the balance between serum transferrin saturation and serum transferrin-receptor concentrations determines which of these functions predominates. The gene which encodes HFE maps to human chromosome 6p22.2.

REFERENCES

1. Cragg, S.J., Drysdale, J. and Worwood, M. 1985. Genes for the "H" sub-unit of human ferritin are present on a number of human chromosomes. *Hum. Genet.* 71: 108-112.
2. McGill, J.R., Naylor, S.L., Sakaguchi, A.Y., Moore, C.M., Boyd, D., Barrett, K.J., Shows, T.B. and Drysdale, J.W. 1987. Human ferritin H and L sequences lie on ten different chromosomes. *Hum. Genet.* 76: 66-72.
3. Waheed, A., Parkkila, S., Saarnio, J., Fleming, R.E., Zhou, X.Y., Tomatsu, S., Britton, R.S., Bacon, B.R. and Sly, W.S. 1999. Association of HFE protein with transferrin receptor in crypt enterocytes of human duodenum. *Proc. Natl. Acad. Sci. USA* 96: 1579-1584.
4. Townsend, A. and Drakesmith, H. 2002. Role of HFE in iron metabolism, hereditary haemochromatosis, anaemia of chronic disease, and secondary iron overload. *Lancet* 359: 786-790.
5. LocusLink Report (LocusID: 235200). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: HFE (human) mapping to 6p22.2.

SOURCE

HFE (W-24) is an affinity purified rabbit polyclonal antibody raised against synthetic HFE peptide of human origin.

PRODUCT

Each vial contains 50 μ g IgG in 500 μ l PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

HFE (W-24) is recommended for detection of HFE of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HFE siRNA (h): sc-43832, HFE shRNA Plasmid (h): sc-43832-SH and HFE shRNA (h) Lentiviral Particles: sc-43832-V.


Molecular Weight of HFE: 49 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227 or human liver extract: sc-363766

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



HFE (W-24): sc-133654. Western blot analysis of HFE expression in Hep G2 whole cell lysate (A) and human liver tissue extract (B).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **HFE (C-12): sc-514405** or **HFE (44-H): sc-130375**, our highly recommended monoclonal alternatives to HFE (W-24).