

IFIT3 (E-22): sc-133687

BACKGROUND

The tetratricopeptide repeat (TPR) motif is a degenerate, 34 amino acid sequence found in many proteins and acts to mediate protein-protein interactions in various pathways. At the sequence level, there can be up to 16 tandem TPR repeats, each of which has a helix-turn-helix shape that stacks on other TPR repeats to achieve ligand binding specificity. IFIT3 (interferon-induced protein with tetratricopeptide repeats 3), also known as IRG2, IFI60, IFIT4, ISG60 or RIG-G, is a 490 amino acid protein that contains eight TPR repeats and may play a role in cell cycle regulation and cellular proliferation. The gene encoding IFIT3 maps to human chromosome 10, which houses over 1,200 genes and comprises nearly 4.5% of the human genome. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie-Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromic deafness, Wolman's syndrome, Cowden syndrome, multiple endocrine neoplasia type 2 and porphyria.

REFERENCES

1. Yu, M., et al. 1997. Cloning of a gene (RIG-G) associated with retinoic acid-induced differentiation of acute promyelocytic leukemia cells and representing a new member of a family of interferon-stimulated genes. *Proc. Natl. Acad. Sci. USA* 94: 7406-7411.
2. Zhu, H., et al. 1997. Use of differential display analysis to assess the effect of human cytomegalovirus infection on the accumulation of cellular RNAs: induction of interferon-responsive RNAs. *Proc. Natl. Acad. Sci. USA* 94: 13985-13990.

CHROMOSOMAL LOCATION

Genetic locus: IFIT3 (human) mapping to 10q23.31.

SOURCE

IFIT3 (E-22) is an affinity purified rabbit polyclonal antibody raised against synthetic IFIT3 peptide of human origin.

PRODUCT

Each vial contains 50 µg IgG in 500 µl PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

APPLICATIONS

IFIT3 (E-22) is recommended for detection of IFIT3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IFIT3 siRNA (h): sc-75326, IFIT3 shRNA Plasmid (h): sc-75326-SH and IFIT3 shRNA (h) Lentiviral Particles: sc-75326-V.


Molecular Weight of IFIT3: 58 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.


DATA




IFIT3 (E-22): sc-133687. Western blot analysis of IFIT3 expression in Caki-1 (A) and COLO 320DM (B) whole cell lysates.



IFIT3 (E-22): sc-133687. Western blot analysis of IFIT3 expression in HeLa whole cell lysate.



IFIT3 (E-22): sc-133687. Western blot analysis of IFIT3 expression in Hep G2 whole cell lysate.



IFIT3 (E-22): sc-133687. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human kidney tissue showing nuclear and cytoplasmic localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **IFIT3 (B-7): sc-393512** or **IFIT3 (E-10): sc-393396**, our highly recommended monoclonal alternatives to IFIT3 (E-22).