

NHEDC1 (K-24): sc-133819

BACKGROUND

The Na⁺/H⁺ exchangers (NHEs) catalyze the transport of Na⁺ in exchange for H⁺ across membranes in organisms and are required for numerous physiological processes. NHEDC1 (Na⁺/H⁺ exchanger domain containing 1), also known as sodium/hydrogen exchanger-like domain-containing protein 1 or NHE domain-containing protein 1, is a 515 amino acid multi-pass membrane protein belonging to the monovalent cation:proton antiporter 1 (CPA1) transporter family. Only expressed in testis, NHEDC1 is highly conserved in mammals including human, mouse, rat and *Macaca fascicularis*. The gene encoding NHEDC1 maps to human chromosome 4, which houses nearly 6% of the human genome and has the largest gene deserts (regions of the genome with no protein encoding genes) of all of the human chromosomes. Defects in some of the genes located on chromosome 4 are associated with Huntington's disease, Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.

REFERENCES

1. Bianchini, L. and Poussegur, J. 1994. Molecular structure and regulation of vertebrate Na⁺/H⁺ exchangers. *J. Exp. Biol.* 196: 337-345.
2. Noël, J. and Pouyssegur, J. 1995. Hormonal regulation, pharmacology, and membrane sorting of vertebrate Na⁺/H⁺ exchanger isoforms. *Am. J. Physiol.* 268: C283-C296.
3. Yun, C.H., Tse, C.M., Nath, S.K., Levine, S.A., Brant, S.R. and Donowitz, M. 1995. Mammalian Na⁺/H⁺ exchanger gene family: structure and function studies. *Am. J. Physiol.* 269: G1-11.
4. Chen, C.S., Bejcek, B.E. and Kersey, J.H. 1995. A mapping study of 13 genes on human chromosome bands 4q11 → q25. *Cytogenet. Cell Genet.* 69: 260-265.
5. Wiebe, C.A., Dibattista, E.R. and Fliegel, L. 2001. Functional role of polar amino acid residues in Na⁺/H⁺ exchangers. *Biochem. J.* 357: 1-10.
6. Ritter, M., Fuerst, J., Wöll, E., Chwatal, S., Gschwentner, M., Lang, F., Deetjen, P. and Paulmichl, M. 2001. Na⁺/H⁺ exchangers: linking osmotic disequilibrium to modified cell function. *Cell. Physiol. Biochem.* 11: 1-18.

CHROMOSOMAL LOCATION

Genetic locus: NHEDC1 (human) mapping to 4q24.

SOURCE

NHEDC1 (K-24) is an affinity purified rabbit polyclonal antibody raised against synthetic NHEDC1 peptide of human origin.

PRODUCT

Each vial contains 50 µg IgG in 500 µl PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

NHEDC1 (K-24) is recommended for detection of NHEDC1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NHEDC1 siRNA (h): sc-89177, NHEDC1 shRNA Plasmid (h): sc-89177-SH and NHEDC1 shRNA (h) Lentiviral Particles: sc-89177-V.


Molecular Weight of NHEDC1: 56 kDa.

Positive Controls: human fetal brain tissue extract.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



NHEDC1 (K-24): sc-133819. Western blot analysis of NHEDC1 expression in human fetal brain tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.