

NNP-1 (P-24): sc-133834

BACKGROUND

Novel nuclear protein 1 (NNP-1), also known as RRP1-like protein or nucleolar protein Nop52, is a 461 amino acid protein belonging to the RRP1 family. Localized to the nucleolus, NNP-1 has simian virus 40-type and bipartite nuclear localization signals and four coiled-coil domains within its C-terminal region. NNP-1 has been found to play an important role in the generation of 28S rRNA in the late processing steps of ribosome biogenesis. At the end of mitosis, nucleolar proteins assemble in a sequential order during the rebuilding of the nucleolus. NNP-1 assembles after Fibrillarin and C23, and simultaneously with B23 and POP1 in the prenucleolar body pathway.

REFERENCES

- Jansen, E., Meulemans, S.M., Orleans, I.C. and Van de Ven, W.J. 1997. The NNP-1 gene (D21S2056E), which encodes a novel nuclear protein, maps in close proximity to the cystatin B gene within the EPM1 and APECED critical region on 21q22.3. *Genomics* 42: 336-341.
- Savino, T.M., Bastos, R., Jansen, E. and Hernandez-Verdun, D. 1999. The nucleolar antigen Nop52, the human homologue of the yeast ribosomal RNA processing RRP1, is recruited at late stages of nucleologenesis. *J. Cell Sci.* 112: 1889-1900.
- Savino, T.M., Gèbrane-Younès, J., De Mey, J., Sibarita, J.B. and Hernandez-Verdun, D. 2001. Nucleolar assembly of the rRNA processing machinery in living cells. *J. Cell Biol.* 153: 1097-1110.
- Scherl, A., Coutè, Y., Dèon, C., Callè, A., Kindbeiter, K., Sanchez, J.C., Greco, A., Hochstrasser, D. and Diaz, J.J. 2002. Functional proteomic analysis of human nucleolus. *Mol. Biol. Cell* 13: 4100-4109.
- Zharskaia, O.O. and Zatspeina, O.V. 2005. Assembly of nucleolus-derived foci in various cultured mammalian cells during mitosis. *Tsitologiya* 47: 780-788.
- Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 610653. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: RRP1 (human) mapping to 21q22.3.

SOURCE

NNP-1 (P-24) is an affinity purified rabbit polyclonal antibody raised against synthetic NNP-1 peptide of human origin.

PRODUCT

Each vial contains 50 µg IgG in 500 µl PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

NNP-1 (P-24) is recommended for detection of NNP-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NNP-1 siRNA (h): sc-91534, NNP-1 shRNA Plasmid (h): sc-91534-SH and NNP-1 shRNA (h) Lentiviral Particles: sc-91534-V.


Molecular Weight of NNP-1: 52 kDa.

Positive Controls: NNP-1 (h): 293T Lysate: sc-176658, Hep G2 cell lysate: sc-2227 or K-562 nuclear extract: sc-2130.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



NNP-1 (P-24): sc-133834. Western blot analysis of NNP-1 expression in non-transfected: sc-117752 (A) and human NNP-1 transfected: sc-176658 (B) 293T whole cell lysates.



NNP-1 (P-24): sc-133834. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human kidney tissue showing nuclear and cytoplasmic localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.