

# NUDT12 (U-25): sc-133855

## BACKGROUND

NUDT12 (nudix (nucleoside diphosphate linked moiety X)-type motif 12) is a 462 amino acid peroxisomal protein belonging to the nudix hydrolase family and the NudC subfamily. Nudix hydrolases contain a characteristic nudix domain and are responsible for catalyzing the hydrolysis of nucleoside diphosphate derivatives. NUDT12 functions as a manganese-dependent polyphosphate phosphohydrolase at an optimum pH of 8-9. Containing three ANK repeats and a nudix hydrolase domain, NUDT12 may act to regulate the concentration of peroxisomal nicotinamide nucleotide cofactors, which are required for oxidative metabolism in peroxisomes. The gene encoding NUDT12 maps to human chromosome 5, which contains 181 million base pairs and comprises nearly 6% of the human genome. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of the q arm or of chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

## REFERENCES

- Dixon, M.J., Read, A.P., Donnai, D., Colley, A., Dixon, J. and Williamson, R. 1991. The gene for Treacher Collins syndrome maps to the long arm of chromosome 5. *Am. J. Hum. Genet.* 49: 17-22.
- Saltman, D.L., Dolganov, G.M., Warrington, J.A., Wasmuth, J.J. and Lovett, M. 1993. A physical map of 15 loci on human chromosome 5q23-q33 by two-color fluorescence *in situ* hybridization. *Genomics* 16: 726-732.
- Abdelraheim, S.R., Spiller, D.G. and McLennan, A.G. 2003. Mammalian NADH diphosphatases of the Nudix family: cloning and characterization of the human peroxisomal NUDT12 protein. *Biochem. J.* 374: 329-335.
- South, S.T., Swensen, J.J., Maxwell, T., Rope, A., Brothman, A.R. and Chen, Z. 2006. A new genomic mechanism leading to Cri du chat syndrome. *Am. J. Med. Genet. A* 140: 2714-2720.
- Du, H.Y., Idol, R., Robledo, S., Ivanovich, J., An, P., Londono-Vallejo, A., Wilson, D.B., Mason, P.J. and Bessler, M. 2007. Telomerase reverse transcriptase haploinsufficiency and telomere length in individuals with 5p- syndrome. *Aging Cell* 6: 689-697.

## CHROMOSOMAL LOCATION

Genetic locus: NUDT12 (human) mapping to 5q21.2; Nudt12 (mouse) mapping to 17 D.

## SOURCE

NUDT12 (U-25) is a Protein A purified rabbit polyclonal antibody raised against synthetic NUDT12 peptide of human origin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

## PRODUCT

Each vial contains 100 µg IgG in 1.0 ml PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

## APPLICATIONS

NUDT12 (U-25) is recommended for detection of NUDT12 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NUDT12 siRNA (h): sc-91746, NUDT12 siRNA (m): sc-150102, NUDT12 shRNA Plasmid (h): sc-91746-SH, NUDT12 shRNA Plasmid (m): sc-150102-SH, NUDT12 shRNA (h) Lentiviral Particles: sc-91746-V and NUDT12 shRNA (m) Lentiviral Particles: sc-150102-V.


Molecular Weight of NUDT12: 52 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

## DATA



NUDT12 (U-25): sc-133855. Western blot analysis of NUDT12 expression in Hep G2 whole cell lysate.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.