

SAP 114 (Z-22): sc-133982

BACKGROUND

SAP 114 (spliceosome associated protein 114), also known as PRP21, PRPF21, SF3A120 or SF3A1, is a 793 amino acid ubiquitously expressed nuclear protein belonging to the SURP protein family and contains two SURP motif repeats and one ubiquitin-like domain. SAP 114 is a subunit of the SF3A splicing factor, a heterotrimeric complex comprised of three subunits that act in tandem to mediate the binding of U2 snRNP to the branchpoint sequence (BPS) in pre-mRNA. The SF3A complex is necessary for the conversion of 15S U2 snRNP into the active 17S protein that is directly involved pre-mRNA splicing events. Functioning as the first subunit of the complex, SAP 114 interacts with subunit 2 (SAP 62) and subunit 3 (SAP 61) via its SURP motifs. SAP 114 is also thought to be involved in the assembly of the E complex, a critical regulator of cell cycle progression from G₁ into S phase in mammalian cells.

REFERENCES

- Chiara, M.D., et al. 1994. Specific protein-protein interactions between the essential mammalian spliceosome-associated proteins SAP 61 and SAP 114. *Proc. Natl. Acad. Sci. USA* 91: 6403-6407.
- Seghezzi, W., et al. 1998. Cyclin E associates with components of the pre-mRNA splicing machinery in mammalian cells. *Mol. Cell. Biol.* 18: 4526-4536.
- Das, R., et al. 2000. Functional association of U2 snRNP with the ATP-independent spliceosomal complex E. *Mol. Cell.* 5: 779-787.
- Will, C.L., et al. 2001. A novel U2 and U11/U12 snRNP protein that associates with the pre-mRNA branch site. *EMBO J.* 20: 4536-4546.
- Online Mendelian Inheritance in Man, OMIM[™]. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 605595. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Jurica, M.S., et al. 2002. Purification and characterization of native spliceosomes suitable for three-dimensional structural analysis. *RNA* 8: 426-439.
- Szabo, A., et al. 2004. Statistical modeling for selecting housekeeper genes. *Genome Biol.* 5: R59.
- Rush, J., et al. 2005. Immunoaffinity profiling of tyrosine phosphorylation in cancer cells. *Nat. Biotechnol.* 23: 94-101.
- Bürkstümmer, T., et al. 2006. An efficient tandem affinity purification procedure for interaction proteomics in mammalian cells. *Nat. Methods* 3: 1013-1019.

CHROMOSOMAL LOCATION

Genetic locus: SF3A1 (human) mapping to 22q12.2; Sf3a1 (mouse) mapping to 11 A1.

SOURCE

SAP 114 (Z-22) is an affinity purified rabbit polyclonal antibody raised against synthetic SAP 114 peptide of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 50 µg IgG in 500 µl PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

APPLICATIONS

SAP 114 (Z-22) is recommended for detection of SAP 114 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SAP 114 siRNA (h): sc-62974, SAP 114 siRNA (m): sc-62975, SAP 114 shRNA Plasmid (h): sc-62974-SH, SAP 114 shRNA Plasmid (m): sc-62975-SH, SAP 114 shRNA (h) Lentiviral Particles: sc-62974-V and SAP 114 shRNA (m) Lentiviral Particles: sc-62975-V.


Molecular Weight of SAP 114: 120 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, 721 B whole cell lysate or K-562 whole cell lysate: sc-2203.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



SAP 114 (Z-22): sc-133982. Western blot analysis of SAP 114 expression in HeLa (A) and K-562 (B) whole cell lysates.



SAP 114 (Z-22): sc-133982. Western blot analysis of SAP 114 expression in 721B whole cell lysate.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.