

SF4 (B-25): sc-133995

BACKGROUND

SF4 (splicing factor 4), also known as RNA-binding protein (RBP) or F23858, is a 645 amino acid nuclear protein and member of the SURP family of splicing factors that is involved in pre-mRNA splicing. Existing as two alternatively spliced isoforms and highly expressed in testis and heart, SF4 is also found in skeletal muscle, brain and kidney where it functions as a component of the spliceosome. SF4 contains two SURP motif repeats, one G-patch domain and is encoded by a gene that maps to human chromosome 19p13.11. Chromosome 19 consists of over 63 million bases, houses approximately 1,400 genes and is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin (Ig) superfamily members, including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family and Fc receptors (FcRs).

REFERENCES

- Olsen, A., Teglund, S., Nelson, D., Gordon, L., Copeland, A., Georgescu, A., Carrano and A., Hammarström, S. 1994. Gene organization of the pregnancy-specific glycoprotein region on human chromosome 19: assembly and analysis of a 700-kb cosmid contig spanning the region. *Genomics* 23: 659-668.
- Teglund, S., Olsen, A., Khan, W.N., Frängsmyr and L., Hammarström, S. 1994. The pregnancy-specific glycoprotein (PSG) gene cluster on human chromosome 19: fine structure of the 11 PSG genes and identification of 6 new genes forming a third subgroup within the carcinoembryonic antigen (CEA) family. *Genomics* 23: 669-684.
- Wang, L., Lin, S.H., Wu, W.G., Kemp, B.L., Walsh, G.L., Hong and W.K., Mao, L. 2000. C-CAM1, a candidate tumor suppressor gene, is abnormally expressed in primary lung cancers. *Clin. Cancer Res.* 6: 2988-2993.
- Trowsdale, J., Barten, R., Haude, A., Stewart, C.A., Beck and S., Wilson, M.J. 2001. The genomic context of natural killer receptor extended gene families. *Immunol. Rev.* 181: 20-38.
- Sampson, N.D., Hewitt, J.E. 2003. SF4 and SFRS14, two related putative splicing factors on human chromosome 19p13.11. *Gene* 305: 91-100.
- Online Mendelian Inheritance in Man, OMIM™. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 607992. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Leeb, T., Müller, M. 2004. Comparative human-mouse-rat sequence analysis of the ICAM gene cluster on HSA 19p13.2 and a 185-kb porcine region from SSC 2q. *Gene* 343: 239-244.

CHROMOSOMAL LOCATION

Genetic locus: SF4 (human) mapping to 19p13.11; Sf4 (mouse) mapping to 8 B3.3.

SOURCE

SF4 (B-25) is an affinity purified rabbit polyclonal antibody raised against synthetic SF4 peptide of human origin.

PRODUCT

Each vial contains 50 µg IgG in 500 µl PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

APPLICATIONS


SF4 (B-25) is recommended for detection of SF4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SF4 siRNA (h): sc-97276, SF4 siRNA (m): sc-153395, SF4 shRNA Plasmid (h): sc-97276-SH, SF4 shRNA Plasmid (m): sc-153395-SH, SF4 shRNA (h) Lentiviral Particles: sc-97276-V and SF4 shRNA (m) Lentiviral Particles: sc-153395-V.

Molecular Weight of SF4: 72 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227.

DATA



SF4 (B-25): sc-133995. Western blot analysis of SF4 expression in Hep G2 whole cell lysate.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **SF4 (B-3): sc-393016** or **SF4 (H-10): sc-514572**, our highly recommended monoclonal alternatives to SF4 (B-25).