



SGLT-3 (D-21): sc-133996

BACKGROUND

Glucose is the main source of energy for mammalian cells and its entry is mediated by various transporters. This process involves seven facilitative (GLUT-1 to -7) and two concentrative (SGLT-1, SGLT-2) glucose transporters and a sensor (SGLT-3). The SGLT family members use the electrochemical gradient of two sodium ions to transport one glucose molecule. The mRNA of SGLTs increase steadily from the fetal period to maturity along with an increase in their functional activity. SGLT-1 is responsible for the uptake of the dietary sugars glucose and galactose from the intestinal lumen, while SGLT-3 is involved in the detection of luminal glucose only. Both the sodium glucose co-transporters SGLT-1 and SGLT-2 are expressed in kidneys. Mutations in the gene encoding SGLT-2 result in familial renal glucosuria (FRG), an isolated disorder of proximal tubular glucose transport, characterized by abnormal urinary glucose excretion in the presence of normal blood glucose levels.

REFERENCES

1. Turk, E., Klisak, I., Bacallao, R., Sparkes, R.S. and Wright, E.M. 1993. Assignment of the human Na⁺/glucose cotransporter gene SGLT-1 to chromosome 22q13.1. *Genomics* 17: 752-754.
2. Yang, Q., Tian, Y., Wada, J., Kashihara, N., Wallner, E., Peterson, D. and Kanwar, Y.S. 2000. Expression characteristics and relevance of sodium glucose cotransporter-1 in mammalian renal tubulogenesis. *Am. J. Physiol. Renal Physiol.* 279: 765-777.
3. Wallner, E.I., Wada, J., Tramonti, G., Lin, S. and Kanwar, Y.S. 2001. Status of glucose transporters in the mammalian kidney and renal development. *Ren. Fail.* 23: 301-310.
4. Stumpel, F., Burcelin, R., Jungermann, K. and Thorens, B. 2001. Normal kinetics of intestinal glucose absorption in the absence of GLUT-2: evidence for a transport pathway requiring glucose phosphorylation and transfer into the endoplasmic reticulum. *PNAS* 98: 11330-11335.
5. Francis, J., Zhang, J., Farhi, A., Carey, H. and Geller, D.S. 2004. A novel SGLT-2 mutation in a patient with autosomal recessive renal glucosuria. *Nephrol. Dial. Transplant.* 19: 2893-2895.
6. Scheepers, A., Joost, H.G. and Schurmann, A. 2004. The glucose transporter families SGLT and GLUT: molecular basis of normal and aberrant function. *J. Parenter. Enteral Nutr.* 28: 364-371.
7. Rahmoune, H., Thompson, P.W., Ward, J.M., Smith, C.D., Hong, G. and Brown, J. 2005. Glucose transporters in human renal proximal tubular cells isolated from the urine of patients with non-Insulin-dependent diabetes. *Diabetes* 54: 3427-3434.
8. Magen, D., Sprecher, E., Zelikovic, I. and Skorecki, K. 2005. A novel missense mutation in SLC5A2 encoding SGLT2 underlies autosomal-recessive renal glucosuria and aminoaciduria. *Kidney Int.* 67: 34-41.

CHROMOSOMAL LOCATION

Genetic locus: SLC5A4 (human) mapping to 22q12.3.

SOURCE

SGLT-3 (D-21) is a Protein A purified rabbit polyclonal antibody raised against synthetic SGLT-3 peptide of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

APPLICATIONS

SGLT-3 (D-21) is recommended for detection of SGLT-3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SGLT-3 siRNA (h): sc-61541, SGLT-3 shRNA Plasmid (h): sc-61541-SH and SGLT-3 shRNA (h) Lentiviral Particles: sc-61541-V.


Molecular Weight of SGLT-3: 72.7 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



SGLT-3 (D-21): sc-133996. Western blot analysis of SGLT-3 expression in Jurkat whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.