SLC43A2 (F-25): sc-134010



The Power to Overtin

BACKGROUND

SLC43A2 (solute carrier family 43, member 2), also known as LAT4 or PP7664, is a 569 amino acid multi-pass membrane protein that belongs to the SLC43A solute transporter family. Expressed in a variety of tissues with highest expression in kidney and placenta, SLC43A2 functions as a sodium- and chloride-independent transport channel protein that facilitates the transport of large, neutral amino acids across membranes. Overexpression of SLC43A2 is associated with head and neck carcinomas, implicating SLC43A2 as a tumor-associated protein. SLC43A2 may be glycosylated and shares 91% sequence similarity with its mouse counterpart, suggesting a conserved function between species. The gene encoding SLC43A2 maps to human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes.

REFERENCES

- Bodoy, S., Martín, L., Zorzano, A., Palacín, M., Estévez, R. and Bertran, J. 2005. Identification of LAT4, a novel amino acid transporter with system L activity. J. Biol. Chem. 280: 12002-12011.
- Zhdanova, N.S. 2007. Comparative mapping of mink (Mustela vison) chromosome 8p: localization of three human BAC clones. Genetika 43: 1074-1078.
- Haase, C., Bergmann, R., Fuechtner, F., Hoepping, A. and Pietzsch, J. 2007. L-type amino acid transporters LAT1 and LAT4 in cancer: uptake of 3-0-methyl-6-18F-fluoro-L-dopa in human adenocarcinoma and squamous cell carcinoma in vitro and in vivo. J. Nucl. Med. 48: 2063-2071.
- 4. Hoffert, J.D., Wang, G., Pisitkun, T., Shen, R.F. and Knepper, M.A. 2007. An automated platform for analysis of phosphoproteomic datasets: application to kidney collecting duct phosphoproteins. J. Proteome Res. 6: 3501-3508.
- Ramadan, T., Camargo, S.M., Herzog, B., Bordin, M., Pos, K.M. and Verrey, F. 2007. Recycling of aromatic amino acids via TAT1 allows efflux of neutral amino acids via LAT2-4F2hc exchanger. Pflugers Arch. 454: 507-516.
- 6. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 610791. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- 7. Vumma, R., Wiesel, F.A., Flyckt, L., Bjerkenstedt, L. and Venizelos, N. 2008. Functional characterization of tyrosine transport in fibroblast cells from healthy controls. Neurosci. Lett. 434: 56-60.

CHROMOSOMAL LOCATION

Genetic locus: SLC43A2 (human) mapping to 17p13.3.

SOURCE

SLC43A2 (F-25) is an affinity purified rabbit polyclonal antibody raised against synthetic SLC43A2 peptide of human origin.

PRODUCT

Each vial contains 50 μg lgG in 500 μl PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

SLC43A2 (F-25) is recommended for detection of SLC43A2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000)

Suitable for use as control antibody for SLC43A2 siRNA (h): sc-94194, SLC43A2 shRNA Plasmid (h): sc-94194-SH and SLC43A2 shRNA (h) Lentiviral Particles: sc-94194-V.


Molecular Weight of SLC43A2 isoforms: 63/53 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



SLC43A2 (F-25): sc-134010. Western blot analysis of SLC43A2 expression in Jurkat whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com