

## FHL-2 (C-16): sc-13409

### BACKGROUND

The four-and-a-half-LIM domain (FHL) proteins include FHL-1 (SLIM1), FHL-2 (SLIM3), FHL-3 (SLIM2) and FHL-4. The signature "half-domain", a single zinc finger domain located in the N-terminal region, differentiates FHLs from other LIM-only proteins, which have numbers of zinc fingers. Specific combinations of FHL proteins elicit selective activation of both CREB and CREM. Skeletal and cardiac muscle express FHL-1 in high levels as compared to the low level of expression in smooth muscle of the colon, small intestine and prostate. FHL-1 localizes to the cytosol of myoblasts, myotubes and differentiated myocytes. FHL-2 is also located in cardiac and skeletal muscle, as well as in placenta and ovary tissues. FHL-3 is found in skeletal muscle, but absent in cardiac muscle. FHL-4 is expressed exclusively by the seminiferous epithelium of the testis, which suggests that FHL-4 is involved in spermatogenesis. The genetic loci for FHLs vary considerably despite similar amino acid sequences among the FHL group.

### REFERENCES

- Morgan, M.J. and Madgwick, A.J. 1996. SLIM defines a novel family of LIM-proteins expressed in skeletal muscle. *Biochem. Biophys. Res. Commun.* 225: 632-638.
- Chan, K.K., et al. 1998. Molecular cloning and characterization of FHL-2, a novel LIM domain protein preferentially expressed in human heart. *Gene* 210: 345-350.
- Lee, S.M., et al. 1998. Chromosomal mapping, tissue distribution and cDNA sequence of four-and-a-half LIM domain protein 1 (FHL-1). *Gene* 216: 163-170.
- Lee, S.M., et al. 1998. Chromosomal mapping of a skeletal muscle specific LIM-only protein FHL-3 to the distal end of the short arm of human chromosome 1. *Somat. Cell Mol. Genet.* 24: 197-202.
- Morgan, M.J. and Madgwick, A.J. 1999. The LIM proteins FHL-1 and FHL-3 are expressed differently in skeletal muscle. *Biochem. Biophys. Res. Commun.* 255: 245-250.

### CHROMOSOMAL LOCATION

Genetic locus: FHL2 (human) mapping to 2q12.1; Fhl2 (mouse) mapping to 1 B.

### SOURCE

FHL-2 (C-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of FHL-2 of human origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-13409 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### APPLICATIONS

FHL-2 (C-16) is recommended for detection of FHL-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


FHL-2 (C-16) is also recommended for detection of FHL-2 in additional species, including equine, canine, bovine and avian.

Suitable for use as control antibody for FHL-2 siRNA (h): sc-37891, FHL-2 siRNA (m): sc-37892, FHL-2 shRNA Plasmid (h): sc-37891-SH, FHL-2 shRNA Plasmid (m): sc-37892-SH, FHL-2 shRNA (h) Lentiviral Particles: sc-37891-V and FHL-2 shRNA (m) Lentiviral Particles: sc-37892-V.

Molecular Weight of FHL-2: 32 kDa.

Positive Controls: FHL-2 (m): 293T Lysate: sc-120254, HeLa nuclear extract: sc-2120 or A-10 nuclear extract: sc-24959.

### DATA



FHL-2 (C-16): sc-13409. Western blot analysis of FHL-2 expression in non-transfected: sc-117752 (A) and mouse FHL-2 transfected: sc-120254 (B) 293T whole cell lysates.

### SELECT PRODUCT CITATIONS

- Yang, Y., et al. 2005. Suppression of FOXO1 activity by FHL-2 through SIRT1-mediated deacetylation. *EMBO J.* 24: 1021-1032.
- Paul, C., et al. 2006. The LIM-only protein FHL-2 is a negative regulator of E4F1. *Oncogene* 25: 5475-5484.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **FHL-2 (F-1): sc-393514** or **FHL-2 (E-8): sc-398866**, our highly recommended monoclonal alternatives to FHL-2 (C-16).