

FEZF2 (W-25): sc-134180

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. FEZF2 (FEZ family zinc finger 2), also known as FEZ, TOF, FEZL, FKSG36 or ZNF312, is a 459 amino acid nuclear protein that belongs to the Krüppel C2H2-type zinc-finger protein family. Considered a transcription repressor, FEZF2 is required for the specification of corticospinal motor neurons and other subcerebral projection neurons. FEZF2 may play a role in layer and neuronal subtype-specific patterning of subcortical projections and axonal fasciculation. FEZF2 controls the development of dendritic arborization and spines of large layer V pyramidal neurons.

REFERENCES

1. Molyneux, B.J., et al. 2007. Molecular development of corticospinal motor neuron circuitry. *Novartis Found. Symp.* 288: 3-15.
2. Leone, D.P., et al. 2008. The determination of projection neuron identity in the developing cerebral cortex. *Curr. Opin. Neurobiol.* 18: 28-35.
3. Russek-Blum, N., et al. 2008. Dopaminergic neuronal cluster size is determined during early forebrain patterning. *Development* 135: 3401-3413.
4. Chen, B., et al. 2008. The FEZF2-CTIP2 genetic pathway regulates the fate choice of subcortical projection neurons in the developing cerebral cortex. *Proc. Natl. Acad. Sci. USA* 105: 11382-11387.
5. Kwan, K.Y., et al. 2008. SOX5 postmitotically regulates migration, postmitotic differentiation, and projections of subplate and deep-layer neocortical neurons. *Proc. Natl. Acad. Sci. USA* 105: 16021-16026.
6. Song, I.S., et al. 2009. Human ZNF312b promotes the progression of gastric cancer by transcriptional activation of the K-Ras gene. *Cancer Res.* 69: 3131-3139.
7. Shimizu, T., et al. 2009. Formation and patterning of the forebrain and olfactory system by zinc-finger genes FEZF1 and FEZF2. *Dev. Growth Differ.* 51: 221-231.
8. Berberoglu, M.A., et al. 2009. FEZF2 expression delineates cells with proliferative potential and expressing markers of neural stem cells in the adult zebrafish brain. *Gene Expr. Patterns* 9: 411-422.
9. Ruby, K.M. and Zheng, B. 2009. Gene targeting in a HUES line of human embryonic stem cells via electroporation. *Stem Cells* 27: 1496-1506.

CHROMOSOMAL LOCATION

Genetic locus: FEZF2 (human) mapping to 3p14.2; Fezf2 (mouse) mapping to 14 A1.

SOURCE

FEZF2 (W-25) is a Protein A purified rabbit polyclonal antibody raised against synthetic FEZF2 peptide of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

APPLICATIONS

FEZF2 (W-25) is recommended for detection of FEZF2 of mouse, rat, human and canine origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FEZF2 siRNA (h): sc-78236, FEZF2 siRNA (m): sc-155683, FEZF2 shRNA Plasmid (h): sc-78236-SH, FEZF2 shRNA Plasmid (m): sc-155683-SH, FEZF2 shRNA (h) Lentiviral Particles: sc-78236-V and FEZF2 shRNA (m) Lentiviral Particles: sc-155683-V.

Molecular Weight of FEZF2 isoforms: 49/31 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.