

ZNF499 (Z-25): sc-134195

BACKGROUND

The BTB (broad-complex, tramtrack and bric a brac) domain, also known as the POZ (poxvirus and zinc finger) domain, is an N-terminal homodimerization domain that contains multiple copies of kelch repeats and/or C₂H₂-type zinc fingers. Proteins that contain BTB domains are thought to be involved in transcriptional regulation via control of chromatin structure and function. ZNF499 (zinc-finger protein 499), also known as ZBTB45 (zinc-finger and BTB domain containing 45), is a 511 amino acid nuclear protein that contains one BTB domain and 4 C₂H₂-type zinc fingers and is thought to be involved in transcriptional regulation events. The gene encoding ZNF499 maps to human chromosome 19, which consists of over 63 million bases, houses approximately 1,400 genes and is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin (Ig) superfamily members, including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family and Fc receptors (Fc Rs).

REFERENCES

- Zimmermann, W., et al. 1988. Chromosomal localization of the carcinoembryonic antigen gene family and differential expression in various tumors. *Cancer Res.* 48: 2550-2554.
- Bardwell, V.J., et al. 1994. The POZ domain: a conserved protein-protein interaction motif. *Genes Dev.* 8: 1664-1677.
- Zollman, S., et al. 1994. The BTB domain, found primarily in zinc-finger proteins, defines an evolutionarily conserved family that includes several developmentally regulated genes in *Drosophila*. *Proc. Natl. Acad. Sci. USA* 91: 10717-10721.
- Ahmad, K.F., et al. 1998. Crystal structure of the BTB domain from PLZF. *Proc. Natl. Acad. Sci. USA* 95: 12123-12128.
- Trettel, F., et al. 2000. A fine physical map of the CACNA1A gene region on 19p13.1-p13.2 chromosome. *Gene* 241: 45-50.
- Buchet-Poyau, K., et al. 2002. Search for the second Peutz-Jeghers syndrome locus: exclusion of the STK13, PRKCG, KLK10, and PSCD2 genes on chromosome 19 and the STK11IP gene on chromosome 2. *Cytogenet. Genome Res.* 97: 171-178.

CHROMOSOMAL LOCATION

Genetic locus: ZBTB45 (human) mapping to 19q13.43.

SOURCE

ZNF499 (Z-25) is an affinity purified rabbit polyclonal antibody raised against synthetic ZNF499 peptide of human origin.

PRODUCT

Each vial contains 50 µg IgG in 500 µl PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

ZNF499 (Z-25) is recommended for detection of ZNF499 of human and canine origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ZNF499 siRNA (h): sc-76988, ZNF499 shRNA Plasmid (h): sc-76988-SH and ZNF499 shRNA (h) Lentiviral Particles: sc-76988-V.


Molecular Weight of ZNF499: 54 kDa.

Positive Controls: A-431 nuclear extract: sc-2122, HeLa nuclear extract: sc-2120 or Jurkat whole cell lysate: sc-2204.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



ZNF499 (Z-25): sc-134195. Western blot analysis of ZNF499 expression in HeLa (A) and A-431 (B) nuclear extracts and human kidney tissue extract (C).



ZNF499 (Z-25): sc-134195. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human kidney tissue showing nuclear localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.