# ZNF621 (B-25): sc-134205



The Power to Question

#### **BACKGROUND**

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. As a member of the krueppel  $C_2H_2$ -type zinc-finger protein family, ZNF621 (Zinc finger protein 621) is a 439 amino acid nuclear protein that contains one KRAB domain and seven  $C_2H_2$ -type zinc fingers. The gene encoding ZNF621 maps to human chromosome 3, which is made up of about 214 million bases encoding over 1,100 genes, including a chemokine receptor (CKR) gene cluster and a variety of human cancer-related gene loci. Marfan Syndrome, porphyria, von Hippel-Lindau syndrome, osteogenesis imperfecta and Charcot-Marie-Tooth Disease are a few of the numerous genetic diseases associated with chromosome 3. There are two isoforms of ZNF621 that are produced as a result of alternative splicing events.

## **REFERENCES**

- Payre, F. and Vincent, A. 1988. Finger proteins and DNA-specific recognition: distinct patterns of conserved amino acids suggest different evolutionary modes. FEBS Lett. 234: 245-250.
- 2. Thiesen, H.J. 1990. Multiple genes encoding zinc finger domains are expressed in human T cells. New Biol. 2: 363-374.
- 3. Rosenfeld, R. and Margalit, H. 1993. Zinc fingers: conserved properties that can distinguish between spurious and actual DNA-binding motifs. J. Biomol. Struct. Dyn. 11: 557-570.
- Müller, S., Stanyon, R., Finelli, P., Archidiacono, N. and Wienberg, J. 2000. Molecular cytogenetic dissection of human chromosomes 3 and 21 evolution. Proc. Natl. Acad. Sci. USA 97: 206-211.
- Braga, E.A., Kashuba, V.I., Maliukova, A.V., Loginov, V.I., Senchenko, V.N., Bazov, I.V., Kiselev, L.L. and Zabarovski, E.R. 2003. New tumor suppressor genes in hot spots of human chromosome 3: new methods of identification. Mol. Biol. 37: 194-211.
- Edelstein, L.C. and Collins, T. 2005. The SCAN domain family of zinc finger transcription factors. Gene 359: 1-17.
- Liu, J. and Stormo, G.D. 2008. Context-dependent DNA recognition code for C<sub>2</sub>H<sub>2</sub> zinc-finger transcription factors. Bioinformatics 24: 1850-1857.
- Ruiz-Herrera, A. and Robinson, T.J. 2008. Evolutionary plasticity and cancer breakpoints in human chromosome 3. Bioessays 30: 1126-1137.

# **CHROMOSOMAL LOCATION**

Genetic locus: ZNF621 (human) mapping to 3p22.1.

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **SOURCE**

ZNF621 (B-25) is a Protein A purified rabbit polyclonal antibody raised against synthetic ZNF621 peptide of human origin.

#### **PRODUCT**

Each vial contains 100  $\mu g$  lgG in 1.0 ml PBS with < 0.1% sodium azide, 0.1% gelatin and < 0.02% sucrose.

#### **APPLICATIONS**

ZNF621 (B-25) is recommended for detection of ZNF621 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ZNF621 siRNA (h): sc-77994, ZNF621 shRNA Plasmid (h): sc-77994-SH and ZNF621 shRNA (h) Lentiviral Particles: sc-77994-V.

Molecular Weight of ZNF621: 49 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204.

#### **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

### **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

**Santa Cruz Biotechnology, Inc.** 1.800.457.3801 831.457.3800 fax 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**