

# podoplanin (M-172): sc-134483

## BACKGROUND

Puromycin aminonucleoside nephrosis (PAN) is a rat model for human minimal change nephropathy. During PAN, severe proteinuria is induced that is paralleled by a reduced expression of a rat podocyte protein, named podoplanin. Podoplanin, also known as glycoprotein 38 (gp38), is a type I membrane protein. Podoplanin localizes in stromal cells of peripheral lymphoid tissue and thymic epithelial cells. As a regulator of the lymphatic endothelium, podoplanin probably plays a role in maintaining the unique shape of podocytes.

## REFERENCES

1. Farr, A.G., et al. 1992. Characterization and cloning of a novel glycoprotein expressed by stromal cells in T-dependent areas of peripheral lymphoid tissues. *J. Exp. Med.* 176: 1477-1482.
2. Farr, A., et al. 1992. Characterization of an antigenic determinant preferentially expressed by type I epithelial cells in the murine thymus. *J. Histochem. Cytochem.* 40: 651-664.
3. Schoppmann, S.F, et al. 2001. Lymphatic microvessel density and lymphovascular invasion assessed by anti-podoplanin immunostaining in human breast cancer. *Anticancer Res.* 21: 2351-2355.
4. Ramirez, M.I., et al. 2003. T1a, a lung type I cell differentiation gene, is required for normal lung cell proliferation and alveolus formation at birth. *Dev. Biol.* 256: 61-72.

## CHROMOSOMAL LOCATION

Genetic locus: Pdpn (mouse) mapping to 4 E1.

## SOURCE

podoplanin (M-172) is a rabbit polyclonal antibody raised against amino acids 1-172 representing full length podoplanin of mouse origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

podoplanin (M-172) is recommended for detection of podoplanin of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for podoplanin siRNA (m): sc-44756, podoplanin shRNA Plasmid (m): sc-44756-SH and podoplanin shRNA (m) Lentiviral Particles: sc-44756-V.


Molecular Weight of podoplanin: 43 kDa.

Positive Controls: mouse kidney extract: sc-2255 or mouse thymus extract: sc-2406.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



podoplanin (M-172): sc-134483. Western blot analysis of podoplanin expression in mouse kidney (A) and mouse lymph node (B) tissue extracts.

## SELECT PRODUCT CITATIONS

1. Song, Y., et al. 2014. Up-regulation of podoplanin involves in neuronal apoptosis in LPS-induced neuroinflammation. *Cell. Mol. Neurobiol.* 34: 839-849.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
Satisfaction  
Guaranteed

Try **podoplanin (8.1.1): sc-53533** or **podoplanin (B-11): sc-166906**, our highly recommended monoclonal alternatives to podoplanin (M-172). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **podoplanin (8.1.1): sc-53533**.