

# ACSL6 (H-70): sc-134498

## BACKGROUND

Acyl-CoA synthetases, also known as long-chain fatty-acid CoA synthases (FACL) or Palmitoyl-CoA ligases, include ACSL1-6, which are all single-pass membrane proteins localizing to the mitochondrion, microsome or peroxisome. ACSL proteins are important for synthesis of cellular lipids and for  $\beta$ -oxidation degradation. Specifically, ACSL proteins catalyze the activation of long-chain fatty acids to acyl-CoAs, which can be metabolized to form CO<sub>2</sub>, triacylglycerol (TAG), phospholipids (PL) and cholesteryl esters (CE). ACSL6 has been shown to be an ETV6 fusion partner gene in a recurrent t(5;12)(q31;p13) translocation in a patient with refractory anemia with excess blasts (RAEB) with basophilia, a patient with acute myelogenous leukemia (AML) with eosinophilia, and a patient with acute eosinophilic leukemia (AEL).

## REFERENCES

1. Yagasaki, F., Jinnai, I., Yoshida, S., Yokoyama, Y., Matsuda, A., Kusumoto, S., Kobayashi, H., Terasaki, H., Ohyashiki, K., Asou, N., Murohashi, I., Bessho, M. and Hirashima, K. 2000. Fusion of TEL/ETV6 to a novel ACS2 in myelodysplastic syndrome and acute myelogenous leukemia with t(5;12)(q31;p13). *Genes Chromosomes Cancer* 26: 192-202.
2. Malhotra, K.T., Malhotra, K., Lubin, B.H. and Kuypers, F.A. 2000. Identification and molecular characterization of acyl-CoA synthetase in human erythrocytes and erythroid precursors. *Biochem. J.* 344: 135-143.
3. Muoio, D.M., Lewin, T.M., Wiedmer, P. and Coleman, R.A. 2001. Acyl-CoAs are functionally channeled in liver: potential role of acyl-CoA synthetase. *Am. J. Physiol. Endocrinol. Metab.* 279: 1366-1373.
4. Coleman, R.A., Lewin, T.M., Van Horn, C.G. and Gonzalez-Baró, M.R. 2002. Do long-chain acyl-CoA synthetases regulate fatty acid entry into synthetic versus degradative pathways? *J. Nutr.* 132: 2123-2126.
5. Qiao, S. and Tuohimaa, P. 2004. The role of long-chain fatty-acid-CoA ligase 3 of prostate cancer LNCaP cell growth. *Biochem. Biophys. Res. Commun.* 319: 358-368.

## CHROMOSOMAL LOCATION

Genetic locus: ACSL6 (human) mapping to 5q31.1; Acs16 (mouse) mapping to 11 B1.3.

## SOURCE

ACSL6 (H-70) is a rabbit polyclonal antibody raised against amino acids 1-70 mapping at the N-terminus of ACSL6 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

ACSL6 (H-70) is recommended for detection of ACSL6 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ACSL6 (H-70) is also recommended for detection of ACSL6 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for ACSL6 siRNA (h): sc-60623, ACSL6 siRNA (m): sc-60624, ACSL6 shRNA Plasmid (h): sc-60623-SH, ACSL6 shRNA Plasmid (m): sc-60624-SH, ACSL6 shRNA (h) Lentiviral Particles: sc-60623-V and ACSL6 shRNA (m) Lentiviral Particles: sc-60624-V.


Molecular Weight of ACSL6: 68-70 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, HEK293 whole cell lysate: sc-45136 or HL-60.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



ACSL6 (H-70): sc-134498. Western blot analysis of ACSL6 expression in Jurkat (A), HEK293 (B), RT-4 (C), HL-60 (D), K-562 (E) and Raji (F) whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.