

KIF4 (H-300): sc-134593

BACKGROUND

The kinesin superfamily proteins (KIFs) are microtubule-dependent molecular motors that transport membranous organelles and protein complexes in a microtubule- and ATP-dependent manner. Cells use KIFs to tightly control the direction, destination and speed of transportation of a variety of important functional molecules, including mRNA. KIF4A functions as an essential chromosome-associated molecular motor involved in faithful chromosome segregation. It is found in the nucleoplasm during interphase and on condensed chromosome arms during mitosis. KIF4A accumulates in the mid-zone during late anaphase and on the cytokinetic ring during cytokinesis. KIF4 binds to and translocates PRC1, a spindle mid-zone-associated cyclin-dependent kinase that plays a role in cytokinesis. KIF4A may also interact with the condensin I and II complexes. Loss of KIF4A leads to chromosome hypercondensation, suggesting that it is necessary for retaining normal chromosome architecture.

REFERENCES

1. Nakagawa, T., et al. 1997. Identification and classification of 16 new kinesin superfamily (KIF) proteins in mouse genome. *Proc. Nat. Acad. Sci. USA* 94: 9654-9659.
2. Miki, H., et al. 2001. All kinesin superfamily protein, KIF, genes in mouse and human. *Proc. Natl. Acad. Sci. USA* 98: 7004-7011.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 300521. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Kurasawa, Y., et al. 2004. Essential roles of KIF4 and its binding partner PRC1 in organized central spindle midzone formation. *EMBO J.* 23: 3237-3248.
5. Mazumdar, M., et al. 2004. Human chromokinesin KIF4A functions in chromosome condensation and segregation. *J. Cell Biol.* 166: 613-620.
6. Zhu, C. and Jiang, W. 2005. Cell cycle-dependent translocation of PRC1 on the spindle by KIF4 is essential for midzone formation and cytokinesis. *Proc. Nat. Acad. Sci. USA* 102: 343-348.

CHROMOSOMAL LOCATION

Genetic locus: KIF4A (human) mapping to Xq13.1, KIF4B (human) mapping to 5q33.2; Kif4 (mouse) mapping to X C3, Kif4b (mouse) mapping to 12 E.

SOURCE

KIF4 (H-300) is a rabbit polyclonal antibody raised against amino acids 361-660 mapping within an internal region of KIF4A of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

KIF4 (H-300) is recommended for detection of KIF4A and KIF4B of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Molecular Weight of KIF4: 140 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Jurkat whole cell lysate: sc-2204 or IMR-32 cell lysate: sc-2409.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



KIF4 (H-300): sc-134593. Western blot analysis of KIF4 expression in DU 145 (A), HeLa (B), Jurkat (C) and IMR-32 (D) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **KIF4 (E-8): sc-365144** or **KIF4 (A-5): sc-365145**, our highly recommended monoclonal alternatives to KIF4 (H-300).