

hCAP-D3 (H-148): sc-134700

BACKGROUND

The condensin complex plays a role in the resolution and segregation of sister chromatids during mitosis and some aspects of mitotic chromosome assembly. Cdc2 phosphorylation of the complex leads to its activation and association with chromosome arms and condensation. Condensin complexes are heteropentamers comprised of two SMC (structural maintenance of chromosomes) subunits and three non-SMC subunits. The SMC family includes SMC1 (also known as SMC1 α and SCMB), which associates with SMC3 (also known as hCAP and Bamacan), SMC2 (also known as hCAP-E), which associates with SMC4 (also known as hCAP-C), and SMC5, which associates with SMC6. hCAP-D3 (non-SMC condensin II complex subunit D3), also known as NCAPD3 or KIAA0056, is a 1498 amino acid nuclear protein that contains 4 HEAT repeats and is a member of the condensin-2 complex.

REFERENCES

1. Nomura, N., Nagase, T., Miyajima, N., Sazuka, T., Tanaka, A., Sato, S., Seki, N., Kawarabayasi, Y., Ishikawa, K. and Tabata, S. 1994. Prediction of the coding sequences of unidentified human genes. II. The coding sequences of 40 new genes (KIAA0041-KIAA0080) deduced by analysis of cDNA clones from human cell line KG-1. *DNA Res.* 1: 223-229.
2. Steen, R.L., Cubizolles, F., Le Guellec, K. and Collas, P. 2000. A kinase-anchoring protein (AKAP)95 recruits human chromosome-associated protein (hCAP)-D2/Eg7 for chromosome condensation in mitotic extract. *J. Cell Biol.* 149: 531-536.
3. Kimura, K., Cuvier, O. and Hirano, T. 2001. Chromosome condensation by a human condensin complex in *Xenopus* egg extracts. *J. Biol. Chem.* 276: 5417-5420.
4. Ball, A.R., Schmiesing, J.A., Zhou, C., Gregson, H.C., Okada, Y., Doi, T. and Yokomori, K. 2002. Identification of a chromosome-targeting domain in the human condensin subunit CNAP1/hCAP-D2/Eg7. *Mol. Cell. Biol.* 22: 5769-5781.
5. Watrin, E. and Legagneux, V. 2005. Contribution of hCAP-D2, a non-SMC subunit of condensin I, to chromosome and chromosomal protein dynamics during mitosis. *Mol. Cell. Biol.* 25: 740-750.

CHROMOSOMAL LOCATION

Genetic locus: NCAPD3 (human) mapping to 11q25; Ncapd3 (mouse) mapping to 9 A4.

SOURCE

hCAP-D3 (H-148) is a rabbit polyclonal antibody raised against amino acids 919-1066 mapping within an internal region of hCAP-D3 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

hCAP-D3 (H-148) is recommended for detection of hCAP-D3 of human origin, Ncapd3 of mouse origin and the corresponding rat homolog by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

hCAP-D3 (H-148) is also recommended for detection of hCAP-D3 in additional species, including equine and canine.

Suitable for use as control antibody for hCAP-D3 siRNA (h): sc-96768, Ncapd3 siRNA (m): sc-149851, hCAP-D3 shRNA Plasmid (h): sc-96768-SH, Ncapd3 shRNA Plasmid (m): sc-149851-SH, hCAP-D3 shRNA (h) Lentiviral Particles: sc-96768-V and Ncapd3 shRNA (m) Lentiviral Particles: sc-149851-V.

Molecular Weight of hCAP-D3: 165 kDa.

Positive Controls: HeLa nuclear extract: sc-2120 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **hCAP-D3 (283.7): sc-101016**, our highly recommended monoclonal alternative to hCAP-D3 (H-148).