Mammaglobin B (H-47): sc-134773



The Power to Question

BACKGROUND

Secretoglobins (Lipophilins, Mammaglobins) are small secreted proteins of endocrine-responsive organs and mucosal epithelia that form multimeric complexes and correlate with the development of various human cancers. Lipophilin A (LIPA, LPHA, secretoglobin, family 1D, member 1; SCGB1D1) and Lipophilin B (LIPB, LPHB, secretoglobin, family 1D, member 2;SCGB1D2) are orthologs of prostatein (estramustine-binding protein), the major secretory glycoprotein of the rat ventral prostate gland. Lipophilin A is a component of a heterodimeric molecule present in human tears. Lipophilin B mRNA can be overexpressed in breast tumors and shows a high degree of correlation with the mRNA expression profile of mammaglobin. Mammaglobin A (MGB1, secretoglobin, family 2A, member 2; SCGB2A2) and Mammaglobin B (MGB2, lipophilin C, LPHC, UGB3, SCGB2A1) histological detection in breast tissue is a reliable diagnostic marker for breast tumors.

REFERENCES

- 1. Wood, D.D., et al. 1984. Interaction between human Myelin basic protein and Lipophilin. Neurochem. Res. 9: 1523-1531.
- Gow, A. 1997. Redefining the Lipophilin family of proteolipid proteins. J Neurosci. Res. 50: 659-664.
- 3. Lehrer, R.I., et al. 1998. Lipophilin, a novel heterodimeric protein of human tears. FEBS Lett. 432: 163-167.
- 4. Carter, D., et al. 2002. Purification and characterization of the Mammaglobin/ Lipophilin B complex, a promising diagnostic marker for breast cancer. Biochemistry 41: 6714-6722.
- Carter, D., et al. 2003. Serum antibodies to Lipophilin B detected in late stage breast cancer patients. Clin. Cancer Res. 9: 749-754.
- Cerveira, N., et al. 2004. Highly sensitive detection of the MGB1 transcript (Mammaglobin) in the peripheral blood of breast cancer patients. Int. J. Cancer 108: 592-595.
- Ouellette, R.J., et al. 2004. RT-PCR for Mammaglobin genes, MGB1 and MGB2, identifies breast cancer micrometastases in sentinel lymph nodes. Am. J. Clin. Pathol. 121: 637-643.
- Sjodin, A., et al. 2005. Secretoglobins in the human pituitary: high expression of Lipophilin B and its downregulation in pituitary adenomas. Acta Neuropathol. 109: 381-386.

CHROMOSOMAL LOCATION

Genetic locus: SCGB2A1 (human) mapping to 11q13.

SOURCE

Mammaglobin B (H-47) is a rabbit polyclonal antibody raised against amino acids 49-95 mapping at the C-terminus of Mammaglobin B of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Mammaglobin B (H-47) is recommended for detection of Mammaglobin B of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Mammaglobin B siRNA (h): sc-45749, Mammaglobin B shRNA Plasmid (h): sc-45749-SH and Mammaglobin B shRNA (h) Lentiviral Particles: sc-45749-V.

Molecular Weight of Mammaglobin B: 9 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com