

# RNH1 (H-135): sc-134824

## BACKGROUND

RNH1, the placental ribonuclease (RNase) inhibitor, is an acidic 460 amino acid protein which contains an unusually high content of leucine and cysteine residues. It is a member of a family of proteinaceous cytoplasmic RNase inhibitors that are expressed in many tissues and bind to both intracellular and extracellular RNases in the cytosol. RNH1 binds to a diverse variety of mammalian RNases and holds them in a latent form. It is also important in the control of mRNA turnover. RNH1 inhibits angiogenesis by reversibly binding angiogenin, a member of the RNaseA superfamily. Because angiogenesis is necessary for the growth and metastasis of tumors, RNH1 may play an important role in cancer gene therapy.

## REFERENCES

- Zhang, B., et al. 2002. Antitumor effect through human endostatin gene transfer in mice bearing B16 melanoma. *Zhonghua Zhong Liu Za Zhi* 24: 451-454.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 173320. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Chen, J.X., et al. 2005. Antitumor effects of human ribonuclease inhibitor gene transfected on B16 melanoma cells. *Int. J. Biochem. Cell Biol.* 37: 1219-1231.
- Dickson, K.A., et al. 2005. Ribonuclease inhibitor: structure and function. *Prog. Nucleic. Acid Res. Mol. Biol.* 80: 349-374.
- Fu, P., et al. 2005. Antitumor effect of hematopoietic cells carrying the gene of ribonuclease inhibitor. *Cancer Gene Ther.* 12: 268-275.
- Iyer, S., et al. 2005. Molecular recognition of human eosinophil-derived neurotoxin (RNase 2) by placental ribonuclease inhibitor. *J. Mol. Biol.* 347: 637-655.
- Rutkoski, T.J., et al. 2005. Disruption of shape-complementarity markers to create cytotoxic variants of ribonuclease A. *J. Mol. Biol.* 354: 41-54.
- Wang, T., et al. 2005. Inhibition of B16 melanoma growth *in vivo* by retroviral vector-mediated human ribonuclease inhibitor. *Angiogenesis* 8: 73-81.

## CHROMOSOMAL LOCATION

Genetic locus: RNH1 (human) mapping to 11p15.5; Rnh1 (mouse) mapping to 7 F5.

## SOURCE

RNH1 (H-135) is a rabbit polyclonal antibody raised against amino acids 1-135 mapping at the N-terminus of RNH1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

RNH1 (H-135) is recommended for detection of RNH1 of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for RNH1 siRNA (h): sc-61365, RNH1 siRNA (m): sc-61366, RNH1 shRNA Plasmid (h): sc-61365-SH, RNH1 shRNA Plasmid (m): sc-61366-SH, RNH1 shRNA (h) Lentiviral Particles: sc-61365-V and RNH1 shRNA (m) Lentiviral Particles: sc-61366-V.


Molecular Weight of RNH1: 50 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227, JAR cell lysate: sc-2276 or K-562 whole cell lysate: sc-2203.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



RNH1 (H-135): sc-134824. Western blot analysis of RNH1 expression in Hep G2 (A), HeLa (B), JAR (C), K-562 (D) and Jurkat (E) whole cell lysates and human liver tissue extract (F).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

**MONOS**  
Satisfaction  
Guaranteed

Try **RNH1 (A-9): sc-365783** or **RNH1 (C-10): sc-271725**, our highly recommended monoclonal alternatives to RNH1 (H-135).