

Nek6 (H-50): sc-134833

BACKGROUND

NIMA was originally shown in *Aspergillus nidulans* to be necessary for entry into mitosis. NIMA-related mammalian proteins have since been identified as Nek1-4 and Nek6-9. High expression of Nek1 is seen in male and female germ cell lines of mice. Nek2 is the closest known mammalian relative to NIMA. Like NIMA, Nek2 expression peaks at the G₂ to M phase transition. Nek3, Nek6, Nek7 and Nek9 also regulate mitosis. Nek1 and Nek8 have been linked with polycystic kidney disease, and Nek4 expression is present in most primary carcinomas. Nek6 localizes to the cytoplasm and is expressed ubiquitously, with highest expression observed in the heart and skeletal muscle. It is activated during M phase and is required for chromosome segregation at the metaphase-anaphase transition and, consequently, mitotic progression.

REFERENCES

- Osmani, S.A., et al. 1988. Mitotic induction and maintenance by overexpression of a G₂-specific gene that encodes a potential protein kinase. *Cell* 53: 237-244.
- Letwin, K., et al. 1992. A mammalian dual specificity protein kinase, Nek1, is related to the NIMA cell cycle regulator and highly expressed in meiotic germ cells. *EMBO J.* 11: 3521-3531.

CHROMOSOMAL LOCATION

Genetic locus: NEK6 (human) mapping to 9q33.3; Nek6 (mouse) mapping to 2 B.

SOURCE

Nek6 (H-50) is a rabbit polyclonal antibody raised against amino acids 1-50 mapping at the N-terminus of Nek6 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Nek6 (H-50) is recommended for detection of Nek6 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Nek6 (H-50) is also recommended for detection of Nek6 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Nek6 siRNA (h): sc-61172, Nek6 siRNA (m): sc-61173, Nek6 shRNA Plasmid (h): sc-61172-SH, Nek6 shRNA Plasmid (m): sc-61173-SH, Nek6 shRNA (h) Lentiviral Particles: sc-61172-V and Nek6 shRNA (m) Lentiviral Particles: sc-61173-V.


Molecular Weight of Nek6: 36 kDa.

Positive Controls: mouse brain extract: sc-2253.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Nek6 (H-50): sc-134833. Western blot analysis of Nek6 expression in mouse brain tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **Nek6 (D-7): sc-374491** or **Nek6 (A-6): sc-393837**, our highly recommended monoclonal alternatives to Nek6 (H-50).