

# GPR128 (M-300): sc-134892

## BACKGROUND

G protein-coupled receptors (GPCRs), also designated seven transmembrane (7TM) receptors and heptahelical receptors, are a protein family which interact with G proteins (heterotrimeric GTPases) to synthesize intracellular second messengers such as diacylglycerol, cyclic AMP, inositol phosphates and calcium ions. Their diverse biological functions range from vision and olfaction to neuronal and endocrine signaling and are involved in many pathological conditions. G protein receptor 128 (GPR128), a member of the secretin family of GPCRs with a GPS domain in its N-terminal domain, may mediate signaling processes to the interior of the cell via activation of G proteins. GPR128 represents an allopeptide which may be involved in T cell mediated transplant rejection as it is able to stimulate 2.102 T cells.

## REFERENCES

1. Lamah, J., Cone, R.I., Maeda, S., Philip, M., Corbani, M., Nádasdi, L., Ramachandran, J., Smith, G.M. and Sadee, W. 1991. Structure and function of G protein coupled receptors. *Pharm. Res.* 7: 1213-1221.
2. Probst, W.C., Snyder, L.A., Schuster, D.I., Brosius, J. and Sealfon, S.C. 1992. Sequence alignment of the G protein-coupled receptor superfamily. *DNA Cell Biol.* 11: 1-20.
3. Fredriksson, R., Gloriam, D.E., Hoglund, P.J., Lagerstrom, M.C. and Schiöth, H.B. 2003. There exist at least 30 human G protein-coupled receptors with long Ser/Thr-rich N-termini. *Biochem. Biophys. Res. Commun.* 301: 752-734.
4. Bjarnadóttir, T.K., Fredriksson, R., Höglund, P.J., Gloriam, D.E., Lagerström, M.C. and Schiöth, H.B. 2004. The human and mouse repertoire of the adhesion family of receptors. *Genomics* 84: 23-33.
5. Lee, J., Hever, A., Willhite, D., Zlotnik, A. and Hevezi, P. 2005. Effects of RNA degradation on gene expression analysis of human postmortem tissues. *FASEB J.* 19: 1356-1358.
6. Felix, N.J., Suri, A., Walters, J.J., Horvath, S., Gross, M.L. and Allen, P.M. 2006. I-Ep-bound self-peptides: identification, characterization, and role in alloreactivity. *J. Immunol.* 176: 1062-1071.

## CHROMOSOMAL LOCATION

Genetic locus: Gpr128 (mouse) mapping to 16 C1.1.

## SOURCE

GPR128 (M-300) is a rabbit polyclonal antibody raised against amino acids 171-470 mapping within an N-terminal extracellular domain of GPR128 of mouse origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

GPR128 (M-300) is recommended for detection of GPR128 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GPR128 siRNA (m): sc-60740, GPR128 shRNA Plasmid (m): sc-60740-SH and GPR128 shRNA (m) Lentiviral Particles: sc-60740-V.

Molecular Weight (predicted) of GPR128: 89 kDa.

Molecular Weight (observed) of GPR128: 84 kDa.

Molecular Weight of glycosylated GPR128: 97 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.