

DCTD (FL-178): sc-134902

BACKGROUND

DCTD (deoxycytidylate deaminase), also known as dCMP deaminase, is a 178 amino acid allosteric enzyme that exists as a homohexamer and belongs to the cytidine and deoxycytidylate deaminase protein family. Using zinc as a cofactor, DCTD catalyzes the deamination of dCMP to dUMP, thereby producing the nucleotide substrate (dUMP) that is used by thymidylate synthase (TS). TS uses 5,10-methyl-enetetrahydrofolate (methylene-THF) and dUMP in the synthesis of 2'-deoxythymidine-5'-monophosphate (dTMP), an essential precursor for DNA biosynthesis. Due to its role in the synthesis of dUMP, DCTD plays an important role in the creation of DNA. The activity of DCTD is regulated by the presence of dCTP and dTTP, two end products in the DCTD metabolic pathway. Multiple isoforms of DCTD are expressed due to alternative splicing events.

REFERENCES

1. Tyrsted, G., et al. 1987. Deoxycytidylate deaminase activity in non-stimulated and phytohemagglutinin-stimulated human lymphocytes, and in leukemic cells. *Mol. Cell. Biochem.* 76: 27-34.
2. Weiner, K.X., et al. 1993. Primary structure of human deoxycytidylate deaminase and overexpression of its functional protein in *Escherichia coli*. *J. Biol. Chem.* 268: 12983-12989.
3. Weiner, K.X., et al. 1995. Chromosomal location and structural organization of the human deoxycytidylate deaminase gene. *J. Biol. Chem.* 270: 18727-18729.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607638. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Rush, J., et al. 2005. Immunoaffinity profiling of tyrosine phosphorylation in cancer cells. *Nat. Biotechnol.* 23: 94-101.
6. Gilbert, J.A., et al. 2006. Gemcitabine pharmacogenomics: cytidine deaminase and deoxycytidylate deaminase gene resequencing and functional genomics. *Clin. Cancer Res.* 12: 1794-1803.
7. Liskay, R.M., et al. 2007. Involvement of deoxycytidylate deaminase in the response to S(n)1-type methylation DNA damage in budding yeast. *Curr. Biol.* 17: R755-R757.
8. Zhang, Y., et al. 2007. Chloroviruses encode a bifunctional dCMP-dCTP deaminase that produces two key intermediates in dTTP formation. *J. Virol.* 81: 7662-7671.
9. Rha, S.Y., et al. 2007. An association between RRM1 haplotype and gemcitabine-induced neutropenia in breast cancer patients. *Oncologist* 12: 622-630.

CHROMOSOMAL LOCATION

Genetic locus: DCTD (human) mapping to 4q35.1; Dctd (mouse) mapping to 8 B1.2.

SOURCE

DCTD (FL-178) is a rabbit polyclonal antibody raised against amino acids 1-178 representing full length DCTD of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

DCTD (FL-178) is recommended for detection of DCTD of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


DCTD (FL-178) is also recommended for detection of DCTD in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for DCTD siRNA (h): sc-89286, DCTD siRNA (m): sc-142907, DCTD shRNA Plasmid (h): sc-89286-SH, DCTD shRNA Plasmid (m): sc-142907-SH, DCTD shRNA (h) Lentiviral Particles: sc-89286-V and DCTD shRNA (m) Lentiviral Particles: sc-142907-V.

Molecular Weight of DCTD: 20 kDa.

Positive Controls: DCTD (h): 293 Lysate: sc-110875, K-562 whole cell lysate: sc-2203 or WI-38 whole cell lysate: sc-364260.

DATA



DCTD (FL-178): sc-134902. Western blot analysis of DCTD expression in non-transfected 293: sc-110760 (A), human DCTD transfected 293: sc-110875 (B), K-562 (C) and WI-38 (D) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **DCTD (F-9): sc-376659**, our highly recommended monoclonal alternative to DCTD (FL-178).