

Pr-Set7 (H-300): sc-135009

BACKGROUND

The methylation of histones plays a pivotal role in the regulation of chromatin structure and gene expression. Histone methylation can occur on Arg or Lys residues, with an exquisite site selectivity for Lys methylation at specific positions in the N-termini of Histones H3 and H4. Pr-Set7, also referred to as SET8, is a nucleosome-specific monomethylase that specifically methylates H4 at Lys 20, a mark of constitutive and facultative heterochromatin. Pr-Set7 is a single subunit enzyme and prefers nucleosomal substrates. It functions to regulate cell-cycle-dependent transcriptional silencing and mitotic regulation in metazoans. The amino acid sequence RHRKVLDRN (17-25) is required for the SET domain of Pr-Set7 to function and, thus, for multiplicity of methylation of Lys 20 of H4 to occur. The methylation mark is very stable and is maintained even in the absence of Pr-Set7.

REFERENCES

1. Fang, J., et al. 2002. Purification and functional characterization of SET8, a nucleosomal Histone H4-Lysine 20-specific methyltransferase. *Curr. Biol.* 12: 1086-1099.
2. Rice, J.C., et al. 2002. Mitotic-specific methylation of Histone H4 Lys 20 follows increased Pr-Set7 expression and its localization to mitotic chromosomes. *Genes Dev.* 16: 2225-2230.

CHROMOSOMAL LOCATION

Genetic locus: SETD8 (human) mapping to 12q24.31; Setd8 (mouse) mapping to 5 F.

SOURCE

Pr-Set7 (H-300) is a rabbit polyclonal antibody raised against amino acids 94-393 mapping at the C-terminus of Pr-Set7 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Pr-Set7 (H-300) is recommended for detection of Pr-Set7 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Pr-Set7 (H-300) is also recommended for detection of Pr-Set7 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Pr-Set7 siRNA (h): sc-62852, Pr-Set7 siRNA (m): sc-155946, Pr-Set7 shRNA Plasmid (h): sc-62852-SH, Pr-Set7 shRNA Plasmid (m): sc-155946-SH, Pr-Set7 shRNA (h) Lentiviral Particles: sc-62852-V and Pr-Set7 shRNA (m) Lentiviral Particles: sc-155946-V.


Molecular Weight of Pr-Set7: 43 kDa.

Positive Controls: NIH/3T3 nuclear extract: sc-2138, RAW 264.7 nuclear extract: sc-24961 or 3611-RF nuclear extract: sc-2143.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Pr-Set7 (H-300): sc-135009. Western blot analysis of Pr-Set7 expression in NIH/3T3 (A), 3611-RF (B), MM-142 (C), RAW 264.7 (D), LADMAC (E) and WEHI-231 (F) nuclear extracts.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Pr-Set7 (D-11): sc-377034** or **Pr-Set7 (D-4): sc-515433**, our highly recommended monoclonal alternatives to Pr-Set7 (H-300).