

Paraplegin (H-180): sc-135026

BACKGROUND

Paraplegin, also known as SPG7 (spastic paraplegia protein 7), CAR, CMAR or PGN, is a 795 amino acid metalloprotease that is a member of the AAA protein family. Localized to the mitochondrial membrane and expressed throughout the body, Paraplegin is a multi-pass membrane protein that is thought to be involved in signal transduction and chaperone-like activities in the mitochondria. Defects in the gene encoding Paraplegin are the cause of spastic paraplegia type 7 (SPG7), a form of autosomal recessive hereditary spastic paraplegia (AR-HSP). HSPs are degenerative spinal cord disorders that are characterized by muscle spasms, stiffness in the legs and, in some cases, incontinence. Recent studies suggest that SPG7 may be a mitochondrial-based disease, as mutations in the Paraplegin gene lead to ragged-red fibers, oxidase-negative fibers and intense succinate dehydrogenase-stained areas of the mitochondria. These mitochondrial dysfunctions lead to axonal degeneration and impaired axonal transport, thus causing the neurodegeneration seen in HSPs.

CHROMOSOMAL LOCATION

Genetic locus: SPG7 (human) mapping to 16q24.3; Spg7 (mouse) mapping to 8 E1.

SOURCE

Paraplegin (H-180) is a rabbit polyclonal antibody raised against amino acids 131-310 mapping within an internal region of Paraplegin of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Paraplegin (H-180) is recommended for detection of Paraplegin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Paraplegin siRNA (h): sc-62755, Paraplegin siRNA (m): sc-62756, Paraplegin shRNA Plasmid (h): sc-62755-SH, Paraplegin shRNA Plasmid (m): sc-62756-SH, Paraplegin shRNA (h) Lentiviral Particles: sc-62755-V and Paraplegin shRNA (m) Lentiviral Particles: sc-62756-V.

Molecular Weight of Paraplegin isoform 1: 88 kDa.


Molecular Weight of Paraplegin isoform 2: 54 kDa.

Positive Controls: Rat skeletal muscle tissue extract.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Paraplegin (H-180): sc-135026. Western blot analysis of Paraplegin expression in rat skeletal muscle tissue extract (A) and human recombinant Paraplegin fusion protein (B).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Paraplegin (C-5): sc-514393**, our highly recommended monoclonal alternative to Paraplegin (H-180).