

COQ7 (FL-217): sc-135040

BACKGROUND

COQ7, a timing protein CLK-1 homolog, is a 217 amino acid protein encoded by the mouse gene *Coq7*. It is believed that COQ7/CLK-1 is required for the biosynthesis of Coenzyme Q (COQ), an essential co-factor in mitochondrial respiration. In yeast, mutation of the COQ7 gene results in the absence of UQ biosynthesis and demonstrates a role for this gene in the step leading to the hydroxylation of 5-demethoxyubiquinone. COQ7 may also be responsible for maintenance of mitochondrial integrity and neurogenesis. COQ7 is highly expressed in tissues with high energy demand such as heart, muscle, liver and kidney. After transcription, COQ7 is targeted to the mitochondria where it is processed to its mature form. The protein similarities and the conservation of function of the CLK-1/CLK-1/COQ7 gene products suggest a potential link between the production of ubiquinone and aging.

CHROMOSOMAL LOCATION

Genetic locus: COQ7 (human) mapping to 16p12.3; *Coq7* (mouse) mapping to 7 F2.

SOURCE

COQ7 (FL-217) is a rabbit polyclonal antibody raised against amino acids 1-217 representing full length COQ7 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

COQ7 (FL-217) is recommended for detection of COQ7 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

COQ7 (FL-217) is also recommended for detection of COQ7 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for COQ7 siRNA (h): sc-62146, COQ7 siRNA (m): sc-62147, COQ7 shRNA Plasmid (h): sc-62146-SH, COQ7 shRNA Plasmid (m): sc-62147-SH, COQ7 shRNA (h) Lentiviral Particles: sc-62146-V and COQ7 shRNA (m) Lentiviral Particles: sc-62147-V.


Molecular Weight of COQ7: 24 kDa.

Positive Controls: mouse heart extract: sc-2254, mouse liver extract: sc-2256 or mouse kidney extract: sc-2255.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



COQ7 (FL-217): sc-135040. Western blot analysis of COQ7 expression in mouse liver (A), mouse heart (B), mouse kidney (C) and rat skeletal muscle (D) tissue extracts.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **COQ7 (F-9): sc-376484** or **COQ7 (B-12): sc-514029**, our highly recommended monoclonal alternatives to COQ7 (FL-217).