CHMP7 (H-300): sc-135050



The Power to Question

BACKGROUND

The charged multivesicular body proteins or chromatin modifying proteins, commonly designated CHMPs, belong to the vacuolar sorting protein family and function as chromatin-modifying proteins. CHMP1-6 are all components of ESCRT (endosomal sorting complex required for transport)-I, -II or -III complexes. These complexes are crucial for sorting endosomal articles into multivesicular bodies (MVBs), as well as required for the formation of these bodies. CHMP7 is an ESCRT-III-related protein that associates with CHMP4B. CHMP7 contains an SNF7 domain in the C-terminus and is most similar to CHMP6. CHMP7 also binds to the microtubule interacting and transport (MIT) domain of the deubiquitinating enzyme, UBPY. The overexpression of CHMP7 inhibits EGFR degradation.

REFERENCES

- Stauffer, D.R., et al. 2001. CHMP1 is a novel nuclear matrix protein affecting chromatin structure and cell-cycle progression. J. Cell Sci. 114: 2383-2393.
- 2. Howard, T.L., et al. 2001. CHMP1 functions as a member of a newly defined family of vesicle trafficking proteins. J. Cell Sci. 114: 2395-2404.
- Yang, K.S., et al. 2004. Molecular characterization of NbCHMP1 encoding a homo-log of human CHMP1 in *Nicotiana benthamiana*. Mol. Cells 17: 255-261.
- Reid, E., et al. 2004. The hereditary spastic paraplegia protein Spastin interacts with the ESCRT-III complex-associated endosomal protein CHMP1B. Hum. Mol. Genet. 14: 19-38.
- 5. Horii, M., et al. 2006. CHMP7, a novel ESCRT-III-related protein, associates with CHMP4B and functions in the endosomal sorting pathway. Biochem. J. 400: 23-32.
- Row, P.E., et al. 2007. The MIT domain of UBPY constitutes a CHMP binding and endosomal localisation signal required for efficient EGF receptor degradation. J. Biol. Chem. 282: 30929-30937.

CHROMOSOMAL LOCATION

Genetic locus: CHMP7 (human) mapping to 8p21.3; Chmp7 (mouse) mapping to 14 D2.

SOURCE

CHMP7 (H-300) is a rabbit polyclonal antibody raised against amino acids 1-300 mapping at the N-terminus of CHMP7 of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

CHMP7 (H-300) is recommended for detection of CHMP7 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CHMP7 (H-300) is also recommended for detection of CHMP7 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for CHMP7 siRNA (h): sc-62106, CHMP7 siRNA (m): sc-62107, CHMP7 shRNA Plasmid (h): sc-62106-SH, CHMP7 shRNA Plasmid (m): sc-62107-SH, CHMP7 shRNA (h) Lentiviral Particles: sc-62106-V and CHMP7 shRNA (m) Lentiviral Particles: sc-62107-V.


Molecular Weight of CHMP7: 51 kDa.

Positive Controls: A549 cell lysate: sc-2413.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CHMP7 (H-300): sc-135050. Western blot analysis of CHMP7 expression in A549 whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **CHMP7 (F-8):** sc-271805, our highly recommended monoclonal alternative to CHMP7 (H-300).