# SANTA CRUZ BIOTECHNOLOGY, INC.

# ALMS1 (H-300): sc-135073



## BACKGROUND

ALMS1 (Alström syndrome protein 1), or ALSS, is a widely expressed protein that localizes to centrosomes and the basal bodies of cilia. It consists of a putative leucine zipper, a tandem repeat domain and a stretch of 17 glutamine residues followed by 7 alanine residues near the N-terminal. Three ALMS1 isoforms exist due to splicing variation. The first isoform is the full length ALMS1. Isoform 2 lacks amino acids 4121-4167. The third isoform is only 3,858 amino acids long and it contains an alternate sequence for amino acids 3,850-3,858. ALMS1 may play a role in intracellular transport, microtubule organization, and basal body and cilia function. A mutation in the gene encoding ALMS1 results in the dysfunction of basal bodies and/or cilia. This dysfunction is suggested to be the cause of Alström syndrome, a rare autosomal-recessive condition. Symptoms include Insulin resistance, type 2 diabetes and obesity.

## REFERENCES

- Tai, T.S., Lin, S.Y. and Sheu, W.H. 2003. Metabolic effects of growth hormone therapy in an Alström syndrome patient. Horm. Res. 60: 297-301.
- Collin, G.B., Cyr, E., Bronson, R., Marshall, J.D., Gifford, E.J., Hicks, W., Murray, S.A., Zheng, Q.Y., Smith, R.S., Nishina, P.M. and Naggert, J.K. 2005. ALMS1-disrupted mice recapitulate human Alström syndrome. Hum. Mol. Genet. 14: 2323-2333.
- Farooqi, I.S. 2005. Genetic and hereditary aspects of childhood obesity. Best Pract. Res. Clin. Endocrinol. Metab. 19: 359-374.
- Hearn, T., Spalluto, C., Phillips, V.J., Renforth, G.L., Copin, N., Hanley, N.A. and Wilson, D.I. 2005. Subcellular localization of ALMS1 supports involvement of centrosome and basal body dysfunction in the pathogenesis of obesity, Insulin resistance, and type 2 diabetes. Diabetes 54: 1581-1587.
- Arsov, T., Larter, C.Z., Nolan, C.J., Petrovsky, N., Goodnow, C.C., Teoh, N.C., Yeh, M.M. and Farrell, G.C. 2006. Adaptive failure to high-fat diet characterizes steatohepatitis in ALMS1 mutant mice. Biochem. Biophys. Res. Commun. 342: 1152-1159.
- Arsov, T., Silva, D.G., O'Bryan, M.K., Sainsbury, A., Lee, N.J., Kennedy, C., Manji, S.S., Nelms, K., Liu, C., Vinuesa, C.G., de Kretser, D.M., Goodnow, C.C. and Petrovsky, N. 2006. Fat aussie—a new Alström syndrome mouse showing a critical role for ALMS1 in obesity, diabetes, and spermatogenesis. Mol. Endocrinol. 20: 1610-1622.

#### CHROMOSOMAL LOCATION

Genetic locus: ALMS1 (human) mapping to 2p13.1.

## SOURCE

ALMS1 (H-300) is a rabbit polyclonal antibody raised against amino acids 2881-3180 mapping within an internal region of ALMS1 of human origin.

# PRODUCT

Each vial contains 200  $\mu g$  lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

ALMS1 (H-300) is recommended for detection of ALMS1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ALMS1 siRNA (h): sc-72345, ALMS1 shRNA Plasmid (h): sc-72345-SH and ALMS1 shRNA (h) Lentiviral Particles: sc-72345-V.


Molecular Weight of ALMS1: 461 kDa.

Positive Controls: human ovary extract: sc-363769.

## **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### DATA



ALMS1 (H-300): sc-135073. Western blot analysis of ALMS1 expression in human ovary tissue extract.

#### **STORAGE**

Store at 4° C, \*\*D0 NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.