

## NNT (H-300): sc-135369

### BACKGROUND

The process of cellular respiration is carried out by integral inner mitochondrial membrane proteins that facilitate the harnessing of energy released by the oxidation of NADH. NNT (nicotinamide nucleotide transhydrogenase), also known as mitochondrial NAD(P) transhydrogenase or pyridine nucleotide transhydrogenase, is a 1,086 amino acid multi-pass mitochondrial inner membrane protein. NNT is a homodimer with an N-terminal section belonging to the AlaDH/PNT family and a C-terminal section belonging to the PNT  $\beta$  subunit family. NNT catalyzes the transfer of a hydride ion from NADH to NADP<sup>+</sup> and functions as a mitochondrial inner membrane proton pump. Using the energy of the proton gradient created by the electron transport chain, NNT produces high concentrations of NADPH, which is used in free radical detoxification and biosynthesis.

### REFERENCES

1. Forsmark-Andrée, P., et al. 1996. Oxidative modification of nicotinamide nucleotide transhydrogenase in submitochondrial particles: effect of endogenous ubiquinol. *Arch. Biochem. Biophys.* 336: 113-120.
2. Arkblad, E.L., et al. 1996. The cDNA sequence of proton-pumping nicotinamide nucleotide transhydrogenase from man and mouse. *Biochim. Biophys. Acta* 1273: 203-205.

### CHROMOSOMAL LOCATION

Genetic locus: NNT (human) mapping to 5p12; Nnt (mouse) mapping to 13 D2.3.

### SOURCE

NNT (H-300) is a rabbit polyclonal antibody raised against amino acids 787-1086 mapping at the C-terminus of NNT of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### APPLICATIONS

NNT (H-300) is recommended for detection of NNT of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NNT (H-300) is also recommended for detection of NNT in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for NNT siRNA (h): sc-91738, NNT siRNA (m): sc-150013, NNT shRNA Plasmid (h): sc-91738-SH, NNT shRNA Plasmid (m): sc-150013-SH, NNT shRNA (h) Lentiviral Particles: sc-91738-V and NNT shRNA (m) Lentiviral Particles: sc-150013-V.


Molecular Weight of NNT: 114 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz<sup>™</sup>: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

### DATA



NNT (H-300): sc-135369. Immunoperoxidase staining of formalin fixed, paraffin-embedded human heart muscle tissue showing cytoplasmic staining of myocytes.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **NNT (B-3): sc-390236** or **NNT (G-8): sc-390215**, our highly recommended monoclonal alternatives to NNT (H-300).