# SANTA CRUZ BIOTECHNOLOGY, INC.

# ARHGAP29 (M-300): sc-135383



## BACKGROUND

GTPase-activating proteins (GAPs) accelerate the intrinsic rate of GTP hydrolysis of Ras-related proteins, resulting in downregulation of their active form. ARHGAP29 (Rho GTPase activating protein 29), also known as PARG1, is a 1,261 amino acid protein that is widely expressed and contains a phorbolester/DAG-type zinc finger and a Rho-GAP domain. There is high expression of ARHGAP29 in skeletal muscle and heart, intermediate expression in placenta, liver and pancreas, and weak expression in brain, lung and kidney. As a GTPase activator, ARHGAP29 converts Rho-type GTPases to an inactive GDP-bound state and has strong activity toward Rho A, and weaker activity toward Rac 1 and Cdc42. Also considered a specific effector of Rap 2A to regulate Rho, ARHGAP29 is strongly down-regulated in mantle-cell lymphomas and upregulated in migrating glioma cells. ARHGAP29 exists as two alternatively spliced isoforms.

## REFERENCES

- Saras, J., et al. 1997. A novel GTPase-activating protein for Rho interacts with a PDZ domain of the protein-tyrosine phosphatase PTPL1. J. Biol. Chem. 272: 24333-24338.
- Bassères, D.S., et al. 2002. ARHGAP10, a novel human gene coding for a potentially cytoskeletal Rho-GTPase activating protein. Biochem. Biophys. Res. Commun. 294: 579-585.
- Myagmar, B.E., et al. 2005. PARG1, a protein-tyrosine phosphatase-associated RhoGAP, as a putative Rap2 effector. Biochem. Biophys. Res. Commun. 329: 1046-1052.
- Meyer-Ficca, M.L., et al. 2005. Poly(ADP-ribose) polymerases: managing genome stability. Int. J. Biochem. Cell Biol. 37: 920-926.
- 5. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 610496. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- Ripperger, T., et al. 2007. Promoter methylation of PARG1, a novel candidate tumor suppressor gene in mantle-cell lymphomas. Haematologica 92: 460-468.

## CHROMOSOMAL LOCATION

Genetic locus: ARHGAP29 (human) mapping to 1p22.1; Arhgap29 (mouse) mapping to 3 G1.

## SOURCE

ARHGAP29 (M-300) is a rabbit polyclonal antibody raised against amino acids 881-1180 mapping within an internal region of ARHGAP29 of mouse origin.

# PRODUCT

Each vial contains 200  $\mu g$  IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

ARHGAP29 (M-300) is recommended for detection of ARHGAP29 of mouse, rat and, to a lesser extent, human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ARHGAP29 siRNA (h): sc-78941, ARHGAP29 siRNA (m): sc-141215, ARHGAP29 shRNA Plasmid (h): sc-78941-SH, ARHGAP29 shRNA Plasmid (m): sc-141215-SH, ARHGAP29 shRNA (h) Lentiviral Particles: sc-78941-V and ARHGAP29 shRNA (m) Lentiviral Particles: sc-141215-V.


Molecular Weight of ARHGAP29: 142 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210.

## **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.





ARHGAP29 (M-300): sc-135383. Western blot analysis of ARHGAP29 expression in HeLa (A) and NIH/3T3 (B) whole cell lysates.

#### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## MONOS Satisfation Guaranteed

Try ARHGAP29 (H-2): sc-377022 or ARHGAP29

(D-3): sc-365554, our highly recommended monoclonal alternatives to ARHGAP29 (M-300).