

TNF β (N-20): sc-1354

BACKGROUND

Tumor necrosis factor β (TNF β), also known as lymphotoxin, is a pleiotropic cytokine that has a molecular weight of 25 kDa. TNF α , also known as cachectin, is a smaller cytokine with a molecular weight of 17 kDa that binds to the same receptors producing a vast array of effects similar to those of TNF β . TNF β and TNF α share 30% amino acid homology and have similar biological activities. TNF β is produced by activated lymphocytes, including CD4⁺ T helper cell type 1 lymphocytes, CD8⁺ lymphocytes and certain B lymphoblastoid cell lines. TNF α is produced by several different cell types, which include lymphocytes, neutrophils and macrophages. TNF α and TNF β can modulate many immune and inflammatory functions, while having the ability to inhibit tumor growth. Target tumor cells must express TNF receptors 1 (55 kDa) and 2 (75 kDa) to be killed, with the p55 receptor mediating the cytotoxic response.

REFERENCES

- Nedwin, G.E., et al. 1985. Human lymphotoxin and tumor necrosis factor genes: structure, homology and chromosomal localization. *Nucleic Acids Res.* 13: 6361-6373.
- Aggarwal, B.B., et al. 1985. Human tumor necrosis factor. Production, purification, and characterization. *J. Biol. Chem.* 260: 2345-2354.
- Vilcek, J., et al. 1991. Tumor necrosis factor. New insights into the molecular mechanisms of its multiple actions. *J. Biol. Chem.* 266: 7313-7316.
- Tartaglia, L.A., et al. 1993. Tumor necrosis factor's cytotoxic activity is signaled by the p55 TNF receptor. *Cell* 73: 213-216.
- De Togni, P., et al. 1994. Abnormal development of peripheral lymphoid organs in mice deficient in lymphotoxin. *Science* 264: 703-707.
- Qin, Z., et al. 1995. Tumor growth inhibition mediated by lymphotoxin: evidence of B lymphocyte involvement in the antitumor response. *Cancer Res.* 55: 4747-4751.
- Sarin, A., et al. 1995. Cytotoxic effect of TNF and lymphotoxin on T lymphoblasts. *J. Immunol.* 155: 3716-3718.

CHROMOSOMAL LOCATION

Genetic locus: LTA (human) mapping to 6p21.3; Lta (mouse) mapping to 17 B1.

SOURCE

TNF β (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of TNF β of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-1354 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

TNF β (N-20) is recommended for detection of TNF β of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


TNF β (N-20) is also recommended for detection of TNF β in additional species, including equine and canine.

Suitable for use as control antibody for TNF β siRNA (h): sc-37218, TNF β siRNA (m): sc-37219, TNF β shRNA Plasmid (h): sc-37218-SH, TNF β shRNA Plasmid (m): sc-37219-SH, TNF β shRNA (h) Lentiviral Particles: sc-37218-V and TNF β shRNA (m) Lentiviral Particles: sc-37219-V.


Molecular Weight of TNF β : 19-25 kDa.

Positive Controls: TNF β (h): 293T Lysate: sc-113691.

DATA



TNF β (N-20): sc-1354. Western blot analysis of TNF β expression in non-transfected: sc-117752 (A) and human TNF β transfected: sc-113691 (B) 293T whole cell lysates.



TNF β (N-20): sc-1354. Western blot analysis of human recombinant TNF β .

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **TNF β (E-6): sc-28345** or **TNF β (D-10): sc-48410**, our highly recommended monoclonal alternatives to TNF β (N-20).