SANTA CRUZ BIOTECHNOLOGY, INC.

HSP 75 (TR1): sc-13557



BACKGROUND

The heat shock proteins (HSPs) comprise a group of highly conserved, abundantly expressed proteins with diverse functions, including the assembly and sequestering of multiprotein complexes, transportation of nascent polypeptide chains across cellular membranes and regulation of protein folding. Heat shock protein 75 mitochondrial precursor (HSP 75), also called tumor necrosis factor type 1 receptor-associated protein (TRAP-1), is a 704 amino acid member of the heat shock protein 90 family. HSP 75 localizes to the mitochondrion. HSP 75 is expressed in a variety of tissues including skeletal muscle, liver, heart, brain, pancreas, lung and placenta and functions as a chaperone that expresses an ATPase activity.

CHROMOSOMAL LOCATION

Genetic locus: TRAP1 (human) mapping to 16p13.3; Trap1 (mouse) mapping to 16 A1.

SOURCE

HSP 75 (TR1) is a mouse monoclonal antibody raised against purified full length HSP 75 of human origin.

PRODUCT

Each vial contains 200 μg lgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

HSP 75 (TR1) is available conjugated to agarose (sc-13557 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-13557 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-13557 PE), fluorescein (sc-13557 FITC), Alexa Fluor[®] 488 (sc-13557 AF488), Alexa Fluor[®] 546 (sc-13557 AF546), Alexa Fluor[®] 594 (sc-13557 AF594) or Alexa Fluor[®] 647 (sc-13557 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-13557 AF680) or Alexa Fluor[®] 790 (sc-13557 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

HSP 75 (TR1) is recommended for detection of HSP 75 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HSP 75 siRNA (h): sc-72191, HSP 75 siRNA (m): sc-72192, HSP 75 shRNA Plasmid (h): sc-72191-SH, HSP 75 shRNA Plasmid (m): sc-72192-SH, HSP 75 shRNA (h) Lentiviral Particles: sc-72191-V and HSP 75 shRNA (m) Lentiviral Particles: sc-72192-V.

Molecular Weight of HSP 75: 75 kDa.

Positive Controls: HL-60 whole cell lysate: sc-2209, Hep G2 cell lysate: sc-2227 or HeLa whole cell lysate: sc-2200.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA





HSP 75 (TR1): sc-13557. Western blot analysis of HSP 75 expression in HL-60 (**A**), Hep G2 (**B**), HeLa (**C**), NCI-H460 (**D**), MCF7 (**E**) and A-673 (**F**) whole cell lysates. Detection reagent used: m-IgG Fc BP-HRP: sr-575409

HSP 75 (TR1): sc-13557. Immunoperoxidase staining of formalin fixed, paraffin-embedded human liver tissue showing cytoplasmic and membrane staining of hepatocytes.

SELECT PRODUCT CITATIONS

- Montesano Gesualdi, N., et al. 2007. Tumor necrosis factor-associated protein 1 (TRAP-1) protects cells from oxidative stress and apoptosis. Stress 10: 342-350.
- Agliarulo, I., et al. 2015. TRAP1 controls cell migration of cancer cells in metabolic stress conditions: correlations with AKT/p70^{S6K} pathways. Biochim. Biophys. Acta 1853: 2570-2579.
- Matassa, D.S., et al. 2016. Oxidative metabolism drives inflammationinduced platinum resistance in human ovarian cancer. Cell Death Differ. 23: 1542-1554.
- Wong, M.Y., et al. 2018. A high-throughput assay for collagen secretion suggests an unanticipated role for HSP 90 in collagen production. Biochemistry 57: 2814-2827.
- Montemurro, L., et al. 2019. A novel MYCN-specific antigene oligonucleotide deregulates mitochondria and inhibits tumor growth in MYCNamplified neuroblastoma. Cancer Res. 79: 6166-6177.
- 6. Criscuolo, D., et al. 2020. Cholesterol homeostasis modulates platinum sensitivity in human ovarian cancer. Cells 9: 828.
- Zhang, L., et al. 2021. TRAP1 inhibits MIC60 ubiquitination to mitigate the injury of cardiomyocytes and protect mitochondria in extracellular acidosis. Cell Death Discov. 7: 389.
- Matassa, D.S., et al. 2022. Regulation of mitochondrial complex III activity and assembly by TRAP1 in cancer cells. Cancer Cell Int. 22: 402.
- Avolio, R., et al. 2023. Cytosolic and mitochondrial translation elongation are coordinated through the molecular chaperone TRAP1 for the synthesis and import of mitochondrial proteins. Genome Res. 33: 1242-1257.

RESEARCH USE

For research use only, not for use in diagnostic procedures.