

MAP-2 (18): sc-135979

BACKGROUND

Microtubules, the primary component of the cytoskeletal network, interact with proteins called microtubule-associated proteins (MAPs). The microtubule-associated proteins can be divided into two groups, structural and dynamic. The structural microtubule-associated proteins, MAP-1A, MAP-1B, MAP-2A, MAP-2B and MAP-2C, stimulate Tubulin assembly, enhance microtubule stability and influence the spatial distribution of microtubules within cells. Both MAP-1 and, to a greater extent, MAP-2 have been implicated as agents of microtubule depolymerization by suppressing the dynamic instability of the microtubules. The suppression of microtubule dynamic instability by the MAP proteins is thought to be associated with phosphorylation of the MAPs.

REFERENCES

1. Sloboda, R.D., et al. 1976. Microtubule-associated proteins and the stimulation of Tubulin assembly *in vitro*. *Biochemistry* 15: 4497-4505.
2. Murphy, D.B., et al. 1977. Role of Tubulin-associated proteins in microtubule nucleation and elongation. *J. Mol. Biol.* 117: 33-52.
3. Hasegawa, M., et al. 1990. Immunochemical evidence that fragments of phosphorylated MAP5 (MAP1B) are bound to neurofibrillary tangles in Alzheimer's disease. *Neuron* 4: 909-918.
4. MacRae, T.H. 1992. Towards an understanding of microtubule function and cell organization: an overview. *Biochem. Cell Biol.* 70: 835-841.
5. Davis, R.J. 1993. The mitogen-activated protein kinase signal transduction pathway. *J. Biol. Chem.* 268: 14553-14556.
6. Maccioni, R.B. and Cambiasso, V. 1995. Role of microtubule-associated proteins in the control of microtubule assembly. *Physiol. Rev.* 75: 835-864.
7. Dhamodharan, R. and Wadsworth, P. 1995. Modulation of microtubule dynamic instability *in vivo* by brain microtubule associated proteins. *J. Cell Sci.* 108: 1679-1689.
8. Vandecandelaere, A., et al. 1996. Differences in the regulation of microtubule dynamics by microtubule-associated proteins MAP1B and MAP2. *Cell Motil. Cytoskeleton* 35: 134-146.

CHROMOSOMAL LOCATION

Genetic locus: MAP2 (human) mapping to 2q34; Mtap2 (mouse) mapping to 1 C3.

SOURCE

MAP-2 (18) is a mouse monoclonal antibody raised against amino acids 19-219 of MAP-2B of human origin.

PRODUCT

Each vial contains 50 µg IgG₁ in 0.5 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

MAP-2 (18) is recommended for detection of MAP-2B of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500); not recommended for immunoprecipitation.


Suitable for use as control antibody for MAP-2 siRNA (h): sc-35853, MAP-2 siRNA (m): sc-35854, MAP-2 shRNA Plasmid (h): sc-35853-SH, MAP-2 shRNA Plasmid (m): sc-35854-SH, MAP-2 shRNA (h) Lentiviral Particles: sc-35853-V and MAP-2 shRNA (m) Lentiviral Particles: sc-35854-V.

Molecular Weight of MAP-2: 280 kDa.


Molecular Weight of MAP-2 low molecular weight isoform: 70 kDa.

Positive Controls: rat brain extract: sc-2392, IMR-32 cell lysate: sc-2409 or SK-N-SH cell lysate: sc-2410.

DATA



MAP-2 (18): sc-135979. Western blot analysis of MAP-2 expression in rat brain tissue extract.



MAP-2 (18): sc-135979. Western blot analysis of MAP-2 expression in IMR-32 whole cell lysate.

SELECT PRODUCT CITATIONS


1. Fan, X.X., et al. 2018. Knockdown of RTN1-C attenuates traumatic neuronal injury through regulating intracellular Ca²⁺ homeostasis. *Neurochem. Int.* 121: 19-25.
2. Filev, A.D., et al. 2019. Oxidized cell-free DNA role in the antioxidant defense mechanisms under stress. *Oxid. Med. Cell. Longev.* 2019: 1245749.
3. Radhakrishnan, S., et al. 2019. Effect of passaging on the stemness of infrapatellar fat pad-derived stem cells and potential role of nucleostemin as a prognostic marker of impaired stemness. *Mol. Med. Rep.* 20: 813-829.

RESEARCH USE

For research use only, not for use in diagnostic procedures. Not for resale.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **MAP-2 (A-4): sc-74421** for MAP-2 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.