SANTA CRUZ BIOTECHNOLOGY, INC.

MEK-3 (30): sc-135985



BACKGROUND

A family of protein kinases located upstream of the MAP kinases and responsible for their activation has been identified. The prototype member of this family, designated MAP kinase kinase, or MEK-1, specifically phosphorylates the MAP kinase regulatory threonine and tyrosine residues present in the Thr-Glu-Tyr motif of ERK. A second MEK family member, MEK-2, resembles MEK-1 in its substrate specificity. MEK-3 (or MKK-3) functions to activate p38 MAP kinase, and MEK-4 (also called SEK1 or MKK-4) activates both p38 and JNK MAP kinases. MEK-5 appears to specifically phosphorylate ERK 5, whereas MEK-6 phosphorylates p38 and p38b. MEK-7 (or MKK-7) phosphorylates and activates the JNK signal transduction pathway.

REFERENCES

- 1. Crews, C.M., et al. 1992. The primary structure of MEK, a protein kinase that phosphorylates the ERK gene product. Science 258: 478-480.
- 2. Wu, J., et al. 1993. Identification and characterization of a new mammalian mitogen-activated protein kinase kinase, MKK-2. Mol. Cell. Biol. 13: 4539-4548.
- 3. Derijard, B., et al. 1995. Independent human MAP-kinase signal transduction pathways defined by MEK and MKK isoforms. Science 267: 682-685.
- 4. Zhou, G., et al. 1995. Components of a new human protein kinase signal transduction pathway. J. Biol. Chem. 270: 12665-12669.
- Han, J., et al. 1996. Characterization of the structure and function of a novel MAP kinase kinase (MKK-6). J. Biol. Chem. 271: 2886-2891.
- 6. Jiang, Y., et al. 1996. Characterization of the structure and function of a new mitogen-activated protein kinase ($p38\beta$). J. Biol. Chem. 271: 17920-17926.
- 7. Tournier, C., et al. 1997. Mitogen-activated protein kinase kinase-7 is an activator of the c-Jun NH_2 -terminal kinase. Proc. Natl. Acad. Sci. USA 94: 7337-7442.
- Holland, P.M., et al. 1997. MKK-7 is a stress-activated mitogen-activated protein kinase kinase functionally related to hemipterous. J. Biol. Chem. 272: 24994-24998.
- Wu, Z., et al. 1997. Molecular cloning and characterization of human JNKK2, a novel Jun NH₂-terminal kinase-specific kinase. Mol. Cell. Biol. 17: 7407-7416.

CHROMOSOMAL LOCATION

Genetic locus: MAP2K3 (human) mapping to 17p11.2; Map2k3 (mouse) mapping to 11 B2.

SOURCE

MEK-3 (30) is a mouse monoclonal antibody raised against amino acids 12-114 of MEK-3b of human origin.

PRODUCT

Each vial contains 50 μg lgG_{2a} in 0.5 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

MEK-3 (30) is recommended for detection of MEK-3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for MEK-3 siRNA (h): sc-35907, MEK-3 siRNA (m): sc-35908, MEK-3 shRNA Plasmid (h): sc-35907-SH, MEK-3 shRNA Plasmid (m): sc-35908-SH, MEK-3 shRNA (h) Lentiviral Particles: sc-35907-V and MEK-3 shRNA (m) Lentiviral Particles: sc-35908-V.

Molecular Weight of MEK-3: 40/37 kDa.

Positive Controls: MEK-3 (h): 293T Lysate: sc-114954, NIH/3T3 whole cell lysate: sc-2210 or Jurkat whole cell lysate: sc-2204.

DATA





MEK-3 (30): sc-135985. Western blot analysis of MEK-3 expression in non-transfected 293T: sc-117752 (**A**), human MEK-3 transfected 293T: sc-114954 (**B**), Jurkat (**C**) and NIH/3T3 (**D**) whole cell lysates. MEK-3 (30): sc-135985. Immunofluorescence staining of methanol-fixed untransfected (**A**) and human MEK-3 transfected HEK 293T cells (**B**).

SELECT PRODUCT CITATIONS

 Ivanov, A.A., et al. 2017. OncoPPi-informed discovery of mitogen-activated protein kinase kinase 3 as a novel binding partner of c-Myc. Oncogene 36: 5852-5860.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures. Not for resale.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.